

Immunoprecipitation Kit – Dynabeads[®] Protein G

For research use only.

INDEX

1. PRODUCT DESCRIPTION
2. PROTOCOLS
3. TECHNICAL ADVICE
4. GENERAL INFORMATION
5. REFERENCES

1. PRODUCT DESCRIPTION

This kit contains Dynabeads[®] Protein G and buffers for immunoprecipitation (see section 1.3 for details).

1.1 Intended Use

This kit is designed for immunoprecipitation of proteins, protein complexes, protein-nucleic acid complexes and other antigens.

1.2 Principle

Antibody (Ab) is added to the Dynabeads Protein G. During a short incubation the Ab will bind to the Dynabeads via their Fc region. The tube is then placed on a Dynal magnet, where the beads will migrate to the side of the tube facing the magnet and allow for easy removal of the supernatant. The bead-bound Ab may now be used for immunoprecipitation. Bound material is easily collected utilizing the unique magnetic properties of the Dynabeads. Magnetic separation facilitates washing, buffer changes and elution (Figure 1).




Fig. 1: Principle of immunoprecipitation of antigen using Dynabeads Protein G.

2. PROTOCOLS

This protocol offers a general guideline for immunoprecipitation. Optimization may be required for each antibody and target antigen (see also section 3). This protocol format uses 50 μ l of Dynabeads Protein G, but may be scaled up or down as required.

2.1 Cell lysis

Cells may be lysed using any standard cell lysis protocol compatible with your starting material. We recommend the use of Cell Extraction Buffer or NP40 Cell Lysis Buffer. For protocols and additional information about cell lysis, please see www.invitrogen.com/immunoprecipitation.

1.3 Description of Materials

Material supplied

- Dynabeads Protein G (2 ml)
These uniform, superparamagnetic beads are 2.8 μ m in diameter, and have recombinant Protein G (approx. 17 kDa) covalently coupled to their surface. The beads are supplied at a concentration of 30 mg/ml in phosphate buffered saline (PBS), pH 7.4, containing 0.01% Tween[®]-20 and 0.09% sodium azide (NaN_3).
- Ab Binding & Washing Buffer (16 ml)
- Washing Buffer (28 ml)
- Elution Buffer (1 ml)

All reagents are of analytical grade. The buffers are compatible with protease/phosphatase inhibitors, which may be used if required.

Additional Materials Required

- Magnet: e.g. DynaMag[™]-2 or DynaMag[™]-Spin (see www.invitrogen.com/magnets for further information and magnet recommendations).
- Mixer allowing rotation of tubes.
- Cell lysis buffer, e.g. Cell Extraction Buffer (Cat. no. FNN0011) or NP40 Cell Lysis Buffer (Cat. no. FNN0021) from Invitrogen
- SDS-PAGE sample buffer, e.g. NuPAGE[®] LDS Sample Buffer (Cat. no. NP0007) and NuPAGE[®] Sample Reducing Agent (Cat. no. NP0009) from Invitrogen

1.4 Antibody selection

The choice of primary antibody is the most important factor for successful target antigen capture. Note that some antibodies may show reduced antigen-binding efficiency for immunoprecipitation, even though the antibody shows good results in other immunological assays. Please refer to the manufacturer's recommendations regarding your primary antibody. See also section 3.

2.2 Preparation of Dynabeads

1. Completely resuspend Dynabeads by pipetting or by rotating on a roller (5 minutes).
2. Transfer 50 μ l (1.5 mg) Dynabeads to a test tube.
3. Separate on the magnet until the supernatant is clear and remove the supernatant.
4. Remove the tube from the magnet.
5. Proceed directly to Ab-binding (section 2.3).

2.3 Binding of Antibody

6. Add your antibody (typically 1-10 μ g Ab), diluted in 200 μ l Ab Binding & Washing Buffer, to the tube from step 4 above. The optimal amount of Ab needed will depend upon the individual Ab used.
7. Incubate with rotation for 10 minutes at room temperature.
8. Place the tube on the magnet and remove the supernatant.
9. Remove the tube from the magnet and resuspend the beads in 200 μ l Ab Binding & Washing Buffer, wash by gentle pipetting.
10. Proceed directly to immunoprecipitation (section 2.4).

Crosslinking

Should you wish to crosslink your antibody to the Dynabeads before continuing with immunoprecipitation, we recommend use of the cross-linking reagent BS³ (5 mM) from Thermo Scientific Group (Pierce). For further information and protocol, please visit: www.invitrogen.com/crosslinking.

2.4 Immunoprecipitation of Target Antigen

11. Place the tube (from step 10) on the magnet, and remove the supernatant.
12. Add your sample containing the antigen (Ag) (typically 100-1,000 μ l) and gently pipette to resuspend the Dynabeads-Ab complex.
13. Incubate with rotation for 10 minutes at room temperature to allow Ag to bind to the Dynabeads-Ab complex. **Note:** Depending on the affinity of the antibody, it may be necessary to increase incubation times for optimal binding. Please see section 3. Technical Advice for details.
14. Place the tube on the magnet. Transfer the supernatant to a clean tube for further analysis, if desired.
15. Wash the Dynabeads-Ab-Ag complex three times using 200 μ l Washing Buffer for each wash. Separate on the magnet between each wash, remove supernatant and resuspend by gentle pipetting.
16. Resuspend the Dynabeads-Ab-Ag complex in 100 μ l Washing Buffer and transfer the bead suspension to a clean tube. This is recommended to avoid co-elution of proteins bound to the tube wall.
17. Proceed to elution (section 2.5, protocol A or B).

2.5 Elution of Target Antigen

Alternative protocols A or B:

A Denaturing elution

18. Place the tube (from step 17) on the magnet and remove the supernatant.
19. Add 20 μ l Elution Buffer and 10 μ l premixed NuPAGE LDS Sample Buffer and NuPAGE Sample Reducing Agent (mixed as per manufacturer's instructions). Gently pipette to resuspend the Dynabeads-Ab-Ag complex.
20. Heat for 10 min at 70°C.
21. Place the tube on the magnet and load the supernatant/sample onto a gel. **Note:** As an alternative, the Dynabeads-Ab-Ag complex can be resuspended in a sample buffer of your choice (e.g. SDS sample buffer). Follow the recommended temperatures and heating times for these buffers prior to gel loading.

B Gentle, non-denaturing elution

18. Place the tube (from step 17) on the magnet and remove the supernatant.
19. Add 20 μ l Elution Buffer and gently pipette to resuspend the Dynabeads-Ab-Ag complex. Avoid foaming.
20. Incubate with rotation for 2 minutes at room temperature to dissociate the complex.
21. Place the tube on the magnet and transfer the supernatant containing eluted Ab and Ag to a clean tube. If the eluted protein is to be used for functional assays or stored, the pH of the eluate can be adjusted by adding 1M Tris, pH 7.5.

3. TECHNICAL ADVICE

Binding characteristics:

Binding capacity:

Dynabeads Protein G have a binding capacity of approximately 7 µg human IgG per mg beads. The amount of Ab captured depends on the concentration of Ab and Dynabeads Protein G in the starting sample.

Antibody affinity:

Ig origin	Affinity for protein A	Affinity for protein G
		A
Human IgG1,2,4	+++	+++
Human IgD	-	-
Human IgA E, M	+	-
Human IgG3	+	+++
Mouse IgG1	+	+++
Mouse IgG2a,2b, 3	+++	+++
Mouse IgM	+	+
Rat IgG1	+	+
Rat IgG2a	-	+++
Rat IgG2b	-	+
Rat IgG2c	+++	+
Bovine IgG1	+	+++
Bovine IgG2	+++	+++
Chicken IgY	-	-
Dog IgG	+++	+
Goat IgG1	+	+++
Goat IgG2	+++	+++
Guinea pig IgG	+++	+
Hamster	+	NA
Horse IgG	+	+++
Monkey IgG	+++	+++
Porcine IgG	+++	+++
Rabbit IgG	+++	+++
Sheep IgG1	+	+++
Sheep IgG2	+++	+++

Table 1: Binding strength of protein A and G to different species of immunoglobulins (Igs) and their subclasses.

Incubation time

Increasing incubation times during immunoprecipitation can improve yield when working with low affinity antibodies. An incubation time of only 10 minutes is sufficient for most applications. Increasing the incubation time to 20-120 minutes can increase yield. Note that non-specific binding can increase with time.

Incubation temperature

For sensitive proteins, to avoid protein complex dissociation and minimize enzymatic activity e.g. when studying phosphorylation events, the isolation protocol including elution (2.5.B) may be run at 4°C.

Storage of Dynabeads-Ab complex

For storage of Ab-conjugated Dynabeads, we recommend using PBS (pH 7.4) with 0.01-0.1 % Tween-20 to prevent aggregation.

Storage of isolated protein

If the immunoprecipitated protein is to be stored, we recommend freezing the Dynabeads-Ab-Ag complex after adding the Elution Buffer and Sample Buffer (step 2.5.A.19). For analysis of the sample, thaw and continue the elution protocol from step. 2.5.A.20.

Technical Support

The latest revision of the package insert/instruction for use is available at www.invitrogen.com.

For further technical information, please visit our website at www.invitrogen.com, or contact Invitrogen Dynal (see www.invitrogen.com/contact for contact details). Certificate of Analysis/Compliance is available upon request.

4. GENERAL INFORMATION

Invitrogen Dynal AS complies with the Quality System Standards ISO 9001:2000 and ISO 13485:2003.

Storage & Stability

Store Dynabeads vials at 2–8°C. Do not freeze the product. Keep Dynabeads in liquid suspension during storage and all handling steps, as drying will result in reduced performance. Resuspend the beads well before use. The buffers may be stored at room temperature. Use care to avoid bacterial/fungal contamination. This product is stable until the expiration date stated on the label, when stored unopened at 2–8°C.

Warnings & Limitations

This product is for research use only. Not intended for any animal or human therapeutic or diagnostic use unless otherwise stated. Preservatives such as sodium azide are toxic if ingested. Avoid pipetting by mouth! Sodium azide may react with lead and copper plumbing to form highly explosive metal azides. When disposing through plumbing drains, flush with large volumes of water to prevent azide build-up. Material Safety Data Sheet (MSDS) is available at <http://www.invitrogen.com>.

Intellectual Property Disclaimer

Invitrogen Dynal will not be responsible for violations or patent infringements that may occur with the use of our products.

Warranty

The products are warranted to the original purchaser only to conform to the quantity and contents stated on the vial and outer labels for the duration of the stated shelf life. Invitrogen Dynal's obligation and the purchaser's exclusive remedy under this warranty is limited either to replacement, at Invitrogen Dynal's expense, of any products which shall be defective in manufacture, and which shall be returned to Invitrogen Dynal, transportation prepaid, or at Invitrogen Dynal's option, refund of the purchase price. Claims for merchandise damaged in transit must be submitted to the carrier.

This warranty shall not apply to any products which shall have been altered outside Invitrogen Dynal, nor shall it apply to any products which have been subjected to misuse or mishandling.

ALL OTHER WARRANTIES, EXPRESSED,

IMPLIED OR STATUTORY, ARE HEREBY SPECIFICALLY EXCLUDED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Invitrogen Dynal's maximum liability is limited in all events to the price of the products sold by Invitrogen Dynal. IN NO EVENT SHALL INVITROGEN DYNAL BE LIABLE FOR ANY SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES. Some states do not allow limits on warranties, or on remedies for breach in certain transactions. In such states, the limits set forth above may not apply.

Limited Use Label License

No. 5: Invitrogen Technology – The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. The buyer may transfer information or materials made through the use of this product to a scientific collaborator, provided that such transfer is not for any Commercial Purpose, and that such collaborator agrees in writing (a) not to transfer such materials to any third party, and (b) to use such transferred materials and/or information solely for research and not for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. Invitrogen Corporation will not assert a claim against the buyer of infringement of patents owned or controlled by Invitrogen Corporation which cover this product based upon the manufacture, use or sale of a therapeutic, clinical diagnostic, vaccine or prophylactic product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. If the purchaser is not willing to accept the limitations of this limited use statement, Invitrogen is willing to accept return of the product with a full refund. For information on purchasing a license to this product for purposes other than research, contact: Licensing Department, Invitrogen Corporation, 1600 Faraday Avenue, Carlsbad, California 92008. Phone (760) 603-7200. Fax (760) 602-6500. Email: outlicensing@invitrogen.com

Trademarks and Patents

Dynal, Dynabeads and DynaMag are either registered trademarks or trademarks of Invitrogen Dynal AS, Oslo, Norway. Any registration or trademark symbols used herein denote the registration status of trademarks in the United States. Trademarks may or may not be registered in other countries.

NuPAGE is a registered trademark of Invitrogen Corporation, USA.

Tween is a registered trademark of ICI Americas, Inc., USA.

Related Dynabeads product

Cat. no.	Name
100.06D	Immunoprecipitation Kit - Dynabeads® Protein A
100.01D/02D	Dynabeads® Protein A
100.03D/04D	Dynabeads® Protein G

A comprehensive range of Dynabeads for use in proteomic workflows are available. Please visit www.invitrogen.com.

5. REFERENCES

Co-immunoprecipitation:

- Lin-Lee Y-C et al. Nuclear localization in the biology of the CD40 receptor in normal and neoplastic human B lymphocytes. *J.Biol.Chem.* 2006;281:18878-18887
- Haren L et al. NEDD1-dependent recruitment of the gamma-tubulin ring complex to the centrosome is necessary for centriole duplication and spindle assembly. *J.Cell Biol.* 2006;172:505-515
- Pham LV et al. Constitutive NF-kappaB and NFAT activation in aggressive B-cell lymphomas synergistically activates the CD154 gene and maintains lymphoma cell survival. *Blood* 2005;106(12):3940-3947
- Catrein I et al. Experimental proof for a signal peptidase I like activity in *Mycoplasma pneumoniae*, but absence of a gene encoding a conserved bacterial type I SPase. *FEBS J.* 2005;272:2892-2900

Purification of IgG from sera:

- Kudva IT et al. Identification of a protein subset of the Anthrax spore immunome in humans immunized with the Anthrax vaccine adsorbed preparation. *Infect. Immun.* 2005;73:5685-5696

Chromatin immunoprecipitation:

- Wu Z et al. Interleukin-21 receptor gene induction in human T cells is mediated by T-cell receptor-induced Sp1 activity. *Mol. Cell.Biol.* 2005;25:9741-9752

Co-immunoprecipitation, automated washing:

- Wan L et al. The survival of motor neurons protein determines the capacity for snRNP assembly: Biochemical deficiency in spinal muscular atrophy. *Mol. Cell. Biol.* 2005;25(13):5543-5551

Invitrogen Dynal is a part of the Invitrogen Group.

Contact details for your local Invitrogen sales office/technical support can be found at <http://www.invitrogen.com/contact>

© Copyright 2008 Invitrogen Dynal AS, Oslo, Norway.

All rights reserved.

SPEC-06597