



Labeled Rabbit Anti-Rat IgG Antibodies

Quick Facts

Storage upon receipt:

• 4°C

Protect from light

Abs/Em: See Table 1

Working Concentration: 1–10 μg/mL

Introduction

Molecular Probes' fluorescent rabbit anti–rat IgG antibodies (Table 1) are prepared from affinity-purified antibodies that react with IgG heavy chains and all classes of immunoglobulin light chains from rat. The Alexa Fluor® dyes to which these antibodies are conjugated provide for extraordinarily bright antibody conjugates. The rabbit anti–rat IgG antibodies have been adsorbed against human IgG prior to labeling to minimize cross-reactivity. The approximate absorption and fluorescence emission maxima for each of the conjugates are shown in Table 1.

In addition to the secondary antibodies described in this Product Information sheet, Molecular Probes prepares fluorescent conjugates of many other species-specific anti-IgG antibodies, as well as conjugates of avidin, streptavidin, NeutrAvidin $^{\rm TM}$ biotin-

Table 1. Molecular Probes' labeled rabbit anti-rat IgG antibodies.*

| Catalog # | Label | Abs † | Em† |
|-----------|-----------------|-------|-----|
| A-21210 | Alexa Fluor 488 | 495 | 519 |
| A-21211 | Alexa Fluor 594 | 590 | 617 |

^{*} These rabbit anti–rat IgG antibodies have been adsorbed against human IgG to minimize cross-reactivity. † Approximate absorption (Abs) and fluorescence emission (Em) maxima in nm.

binding protein, protein A and protein G. Please consult our Web site at www.probes.com or contact our Technical Assistance Department for more information about these products.

Materials

Contents

The fluorophore-labeled rabbit anti-rat IgG (H+L) antibodies are supplied in unit sizes of 0.5 mL as 2 mg/mL solutions in 0.1 M sodium phosphate, 0.1 M NaCl, pH 7.5, containing 5 mM sodium azide.

The degree of labeling for each conjugate is typically 2–8 fluorophore molecules per IgG molecule; the exact degree of labeling is indicated on the product label. At the time of preparation, the products are certified to be free of unconjugated dyes and are tested in a cytological experiment to ensure low nonspecific staining.

Storage

When these products are stored undiluted at 4°C and protected from light, they are stable for at least three months. For longer storage, divide the solution into single-use aliquots and freeze at -20°C. Frozen aliquots are stable for at least six months. PROTECT FROM LIGHT. AVOID REPEATED FREEZING AND THAWING.

Application

It is a good practice to centrifuge the protein conjugate solution briefly in a microcentrifuge before use; only the supernatant should then be added to the experiment. This step will eliminate any protein aggregates that may have formed during storage, thereby reducing nonspecific background staining.

Because staining protocols vary with application, the appropriate dilution of antibody should be determined empirically. For fluorophore-labeled antibodies, a final concentration of $1{\text -}10~\mu\text{g/mL}$ should be satisfactory for most immunohistochemical applications. 1

References

1. Short Protocols in Molecular Biology, 2nd Edition, F.M. Ausubel et al., Eds., John Wiley and Sons (1992) pp. 14-24-14-30.

Product List Current prices may be obtained from our Web site or from our Customer Service Department.

| Cat # | Product Name | Unit Size |
|---------|--|-----------|
| A-21210 | Alexa Fluor® 488 rabbit anti-rat IgG (H+L) conjugate *2 mg/mL* | 0.5 mL |
| A-21211 | Alexa Fluor® 594 rabbit anti-rat IgG (H+L) conjugate *2 mg/mL* | 0.5 mL |

Contact Information

Further information on Molecular Probes' products, including product bibliographies, is available from your local distributor or directly from Molecular Probes. Customers in Europe, Africa and the Middle East should contact our office in Leiden, the Netherlands. All others should contact our Technical Assistance Department in Eugene, Oregon.

Please visit our Web site — www.probes.com — for the most up-to-date information

Molecular Probes, Inc.

PO Box 22010, Eugene, OR 97402-0469 Phone: (541) 465-8300 • Fax: (541) 344-6504

Customer Service: 7:00 am to 5:00 pm (Pacific Time)

Phone: (541) 465-8338 • Fax: (541) 344-6504 • order@probes.com

Toll-Free Ordering for USA and Canada:

Order Phone: (800) 438-2209 • Order Fax: (800) 438-0228

Technical Assistance: 8:00 am to 4:00 pm (Pacific Time)

Phone: (541) 465-8353 • Fax: (541) 465-4593 • tech@probes.com

Molecular Probes Europe BV

PoortGebouw, Rijnsburgerweg 10 2333 AA Leiden, The Netherlands

Phone: +31-71-5233378 • Fax: +31-71-5233419

Customer Service: 9:00 to 16:30 (Central European Time)

Phone: +31-71-5236850 • Fax: +31-71-5233419

eurorder@probes.nl

Technical Assistance: 9:00 to 16:30 (Central European Time)

Phone: +31-71-5233431 • Fax: +31-71-5241883

eurotech@probes.nl

Molecular Probes' products are high-quality reagents and materials intended for research purposes only. These products must be used by, or directly under the supervision of, a technically qualified individual experienced in handling potentially hazardous chemicals. Please read the Material Safety Data Sheet provided for each product; other regulatory considerations may apply.

Several of Molecular Probes' products and product applications are covered by U.S. and foreign patents and patents pending. Our products are not available for resale or other commercial uses without a specific agreement from Molecular Probes, Inc. We welcome inquiries about licensing the use of our dyes, trademarks or technologies. Please submit inquiries by e-mail to busdev@probes.com. All names containing the designation ® are registered with the U.S. Patent and Trademark Office.

Copyright 2001, Molecular Probes, Inc. All rights reserved. This information is subject to change without notice.