

## PRODUCT INFORMATION

# XhoI

**#ER0691**      2000 u

**Lot:**                      **Expiry Date:**

5'...**C↓T** C G A G...3'

3'...**G A G C T↑C**...5'

Concentration:      10 u/μl  
Supplied with:      1 ml of 10X Buffer R  
                                 1 ml of 10X Buffer Tango

**Store at -20°C**



In total 3 vials.

BSA included

[www.thermoscientific.com/fermentas](http://www.thermoscientific.com/fermentas)

## RECOMMENDATIONS

**1X Buffer R** (for 100% XhoI digestion)

10 mM Tris-HCl (pH 8.5), 10 mM MgCl<sub>2</sub>, 100 mM KCl, 0.1 mg/ml BSA.

**Incubation temperature**

37°C.

**Unit Definition**

One unit is defined as the amount of XhoI required to digest 1 μg of lambda DNA-HindIII fragments in 1 hour at 37°C in 50 μl of recommended reaction buffer.

**Dilution**

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl (pH 7.4 at 25°C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/ml BSA and 50% glycerol.

**Double Digests**

Thermo Scientific Tango Buffer is provided to simplify buffer selection for double digests. 98% of Thermo Scientific restriction enzymes are active in a 1X or 2X concentration of Tango™ Buffer. Please refer to to [www.fermentas.com/doubledigest](http://www.fermentas.com/doubledigest) to choose the best buffer for your experiments.

1X Tango Buffer: 33 mM Tris-acetate (pH 7.9 at 37°C), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/ml BSA.

**Storage Buffer**

XhoI is supplied in: 10 mM Tris-HCl (pH 7.4 at 25°C), 100 mM KCl, 1 mM DTT, 1 mM EDTA, 0.2 mg/ml BSA and 50% glycerol.

## Recommended Protocol for Digestion

- Add:

nuclease-free water	16 $\mu$ l
10X Buffer R	2 $\mu$ l
DNA (0.5-1 $\mu$ g/ $\mu$ l)	1 $\mu$ l
XhoI	0.5-2 $\mu$ l*
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

The digestion reaction may be scaled either up or down.

## Recommended Protocol for Digestion of PCR Products Directly after Amplification

- Add:

PCR reaction mixture	10 $\mu$ l (~0.1-0.5 $\mu$ g of DNA)
nuclease-free water	18 $\mu$ l
10X Buffer R	2 $\mu$ l
XhoI	1-2 $\mu$ l*
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

\* This volume of the enzyme is recommended for preparations of standard concentrations (10 u/ $\mu$ l), whereas HC enzymes (50 u/ $\mu$ l) should be diluted with Dilution Buffer to obtain 10 u/ $\mu$ l concentration.

## Thermal Inactivation

XhoI is inactivated by incubation at 80°C for 20 min.

## ENZYME PROPERTIES

### Enzyme Activity in Thermo Scientific REase Buffers, %

B	G	O	R	Tango	2X Tango
0-20	50-100	50-100	100	20-50	100

### Methylation Effects on Digestion

Dam: never overlaps – no effect.

Dcm: never overlaps – no effect.

CpG: completely overlaps – cleavage impaired.

EcoKI: never overlaps – no effect.

EcoBI: never overlaps – no effect.

### Stability during Prolonged Incubation

A minimum of 0.1 units of the enzyme is required for complete digestion of 1  $\mu$ g of lambda DNA in 16 hours at 37°C.

### Digestion of Agarose-embedded DNA

A minimum of 5 units of the enzyme is required for complete digestion of 1  $\mu$ g of agarose-embedded lambda DNA in 16 hours.

### Compatible Ends

Eco88I, Sall, SmoI, SgrDI.

### Number of Recognition Sites in DNA

$\lambda$	$\Phi$ X174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
1	1	0	0	0	0	0

### Note

Supercoiled plasmids may require up to 5-fold more XhoI for complete digestion than linear DNA.

# CERTIFICATE OF ANALYSIS

## Overdigestion Assay

No detectable change in the specific fragmentation pattern is observed after a 160-fold overdigestion with XhoI (10 u/μg lambda DNA x 16 hours).

## Ligation/Recutting Assay

After a 50-fold overdigestion (3 u/μg ΦX174 DNA x 17 hours) with XhoI, more than 95% of the digested DNA fragments can be ligated at a 5'-termini concentration of 0.08 μM. More than 95% of these sites can be recut.

## Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded or double-stranded labeled oligonucleotides occurred during incubation with 10 units of XhoI for 4 hours.

## Blue/White Cloning Assay

A mixture of pUC57/HindIII, pUC57/PstI and pUC57/Eco32I digests was incubated with 10 units of XhoI for 16 hours. After religation and transformation, the background level of white colonies was <1%.

Quality authorized by:

 Jurgita Zilinskiene

## **PRODUCT USE LIMITATION**

This product is developed, designed and sold exclusively *for research purposes and in vitro use only*. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to [www.thermoscientific.com/fermentas](http://www.thermoscientific.com/fermentas) for Material Safety Data Sheet of the product.

© 2011 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.

