

PRODUCT INFORMATION

BclI

#ER0721 1000 u

Lot: **Expiry Date:**

5'...**T↓G A T C A**...3'

3'...**A C T A G↑T**...5'

Concentration: 10 u/μl
Source: *Bacillus caldolyticus*
Supplied with: 1 ml of 10X Buffer G
1 ml of 10X Buffer Tango

Store at -20°C



In total 3 vials.

BSA included

RECOMMENDATIONS

1X Buffer G (for 100% BclI digestion)

10 mM Tris-HCl (pH 7.5), 10 mM MgCl₂, 50 mM NaCl,
0.1 mg/ml BSA.

Incubation temperature

55°C*.

Unit Definition

One unit is defined as the amount of BclI required to digest 1 μg of lambda DNA *dam*⁻ in 1 hour at 55°C in 50 μl of recommended reaction buffer.

Dilution

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl (pH 7.4 at 25°C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/ml BSA and 50% glycerol.

Double Digests

Thermo Scientific Tango Buffer is provided to simplify buffer selection for double digests. 98% of Thermo Scientific restriction enzymes are active in a 1X or 2X concentration of Tango™ Buffer. Please refer to to www.fermentas.com/doubledigest to choose the best buffer for your experiments.

1X Tango Buffer: 33 mM Tris-acetate (pH 7.9 at 37°C), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/ml BSA.

* Incubation at 37°C results in 50% activity.

Storage Buffer

BclI is supplied in: 10 mM Tris-HCl (pH 7.4 at 25°C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/ml BSA and 50% glycerol.

Recommended Protocol for Digestion

- Add:

nuclease-free water	16 μ l
10X Buffer G	2 μ l
DNA (0.5-1 μ g/ μ l)	1 μ l
BclI	0.5-2 μ l**
- Mix gently and spin down for a few seconds.
- Incubate at 55°C for 1-16 hours**.

The digestion reaction may be scaled either up or down.

Recommended Protocol for Digestion of PCR Products Directly after Amplification

- Add:

PCR reaction mixture	10 μ l (~0.1-0.5 μ g of DNA)
nuclease-free water	18 μ l
10X Buffer G	2 μ l
BclI	1-2 μ l**
- Mix gently and spin down for a few seconds.
- Incubate at 55°C for 1-16 hours**.

Thermal Inactivation

Only small amounts of BclI (up to 10 units) can be inactivated at 80°C in 20 min.

Inactivation Procedure

- To prepare the digested DNA for electrophoresis:
 - stop the digestion reaction by adding 0.5M EDTA, pH 8.0 (#R1021), to achieve a 20 mM final concentration. Mix thoroughly, add an electrophoresis loading dye and load onto gel.
- To prepare DNA suitable for further enzymatic reactions:
 - extract with phenol/chloroform, precipitate with ethanol or isopropanol, wash the pellet with 75% cold ethanol and air-dry;
 - dissolve DNA in either nuclease-free water, TE buffer, or a buffer suitable for further applications;
 - check the DNA concentration in the solution.

For **ENZYME PROPERTIES** and **CERTIFICATE OF ANALYSIS**
see back page

** See Star Activity on back page.

ENZYME PROPERTIES

Enzyme Activity in Thermo Scientific REase Buffers, %

B	G	O	R	Tango	2X Tango
20-50	100	20-50	20-50	100***	100

***Star activity appears at a greater than 5-fold overdigestion (5 u x 1h).

Star Activity

An excess of BclI (20 u/μg DNA x 1 hour) may result in star activity.

Methylation Effects on Digestion

Dam: completely overlaps – blocked.

Dcm: never overlaps – no effect.

CpG: never overlaps – no effect.

EcoKI: never overlaps – no effect.

EcoBI: may overlap – blocked.

Stability during Prolonged Incubation

A minimum of 0.1 units of enzyme is required for complete digestion of 1 μg of lambda DNA in 16 hours at 55°C.

Compatible Ends

BamHI, BglII, Bsp143I, MboI, PstI

Number of Recognition Sites in DNA

λ	ΦX174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
8	0	0	0	0	0	0

Note

BclI is blocked by overlapping *dam* methylation. To avoid *dam* methylation, use a *dam*⁻, *dcm*⁻ strain such as GM2163 (#M0099).

CERTIFICATE OF ANALYSIS

Overdigestion Assay

No detectable change in the specific fragmentation pattern is observed after a 15-fold overdigestion with BclI (15 u/μg lambda DNA *dam*⁻ x 1 hour) (see Star Activity).

Ligation/Recutting Assay

After a 10-fold overdigestion (5 u/μg DNA x 2 hours) with BclI, more than 95% of the digested DNA fragments can be ligated at a 5'-termini concentration of 0.1 μM. More than 95% of these sites can be recut.

Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded or double-stranded labeled oligonucleotides occurred during incubation with 10 units of BclI for 4 hours.

Blue/White Cloning Assay

A mixture of pUC57/HindIII, pUC57/PstI and pUC57/Eco32I digests was incubated with 10 units of BclI for 1 hour. After religation and transformation, the background level of white colonies was <1%.

Quality authorized by:



Jurgita Zilinskiene

PRODUCT USE LIMITATION

This product is developed, designed and sold exclusively for research purposes and *in vitro* use only. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to www.thermoscientific.com/fermentas for Material Safety Data Sheet of the product.

© 2011 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.

