

Nuclear Yellow [Hoechst S769121]


Ordering Information

Product Numbers: 17539

Storage Conditions

Avoid exposure to light
Keep at -20 °C and desiccated
Expiration: 24 months upon receipt

Chemical and Physical Properties



Molecular Weight: 651.01

Appearance: Light yellow powder

Solvents: water or dimethylsulfoxide (DMSO)

Spectral Properties: Excitation = 355 nm; Emission = 495 nm.

Biological Applications

The Hoechst stains are a family of fluorescent stains for labeling DNA used in fluorescence microscopy and flow cytometry. Because these fluorescent stains label DNA, they are also commonly used to visualize nuclei and mitochondria. Two of these closely related bis-benzimidazoles, Hoechst 33258 and Hoechst 33342, are commonly used. Both dyes are excited by ultraviolet light at around 350 nm, and both emit blue/cyan fluorescence light with an emission maximum at 461 nm. The Hoechst stains may be used with live or fixed cells, and are often used as a substitute for another nucleic acid stain, DAPI. The key difference between them is that the additional ethyl group of Hoechst 33342 makes it more lipophilic, and thus more readily to cross intact cell membranes. In some applications, Hoechst 33258 is significantly less permeant than Hoechst 33342. These dyes can also be used to detect the contents of a sample DNA by plotting a standard emission-to-content curve.

The long-wavelength tracer nuclear yellow is often combined with the popular retrograde tracer True Blue for two-color neuronal mapping. In neuronal cells, nuclear yellow primarily stains the nucleus with yellow fluorescence, whereas true blue is a UV light-excitable, divalent cationic dye that stains the cytoplasm with blue fluorescence. Both Nuclear Yellow and True Blue are stable when subjected to immunohistochemical processing and can be used to photoconvert DAB into an insoluble, electron-dense reaction product. Hoechst 33258, Hoechst 33342 and Nuclear Yellow, when transported retrogradely through axons to their parent cell bodies, may migrate out of the axons and the cell bodies, as indicated by fluorescence of adjoining glial nuclei. This migration was found to take place both in vivo and in vitro during storage of the sections in water. When using the tracers in 1% concentrations the in vivo migration may be controlled by restricting the survival time. The in vitro migration may be prevented by rapid histological processing of the material.

Sample Protocol for Staining Cells

The following procedure can be adapted for most cell types. Growth medium, cell density, the presence of other cell types and other factors may influence staining. Residual detergent on glassware may also affect real or apparent staining of many organisms, causing brightly stained material to appear in solutions with or without cells present.

- 1) Pellet cells by centrifugation
- 2) Resuspend the cells in buffered solutions or media with optimal dye binding at pH 7.4.
- 3) Stain adherent cells *in situ* on cover slips or in the cell culture wells.
- 4) Add Hoechst stain using the concentrations between 0.5 and 5 μ M.
- 5) Incubate the cells for 15 to 60 minutes.

Note: The incubation time and dye concentration need experimentally tested. In initial experiments, it may be best to try several dye concentrations over the entire suggested range to determine the concentration that yields optimal staining.

References

1. Pinna-Senn E, Lisanti JA, Ortiz MI, Dalmaso G, Bella JL, Gosalvez J, Stockert JC. (2000) Specific heterochromatic banding of metaphase chromosomes using nuclear yellow. *Biotech Histochem*, 75, 132.
2. Santos ML, Lu E, Wolfe J. (2000) Nuclear death in living Tetrahymena: the case of the haploid nuclei. *J Eukaryot Microbiol*, 47, 493.
3. Oda Y. (1986) Nerve center of the rat extraocular muscles as studied by fluorescent dyes. II: Double labeling study with fast blue and nuclear yellow. *Okajimas Folia Anat Jpn*, 63, 141.
4. Fryszak T, Zenker W, Kantner D. (1984) Afferent and efferent innervation of the rat esophagus. A tracing study with horseradish peroxidase and nuclear yellow. *Anat Embryol (Berl)*, 170, 63.
5. Stanzani S, Russo A, Leanza G. (1984) [Projections to the supraoptic and paraventricular nuclei studied in the rat by means of retrograde fluorescent tracers (fast blue, nuclear yellow)]. *Boll Soc Ital Biol Sper*, 60, 2339.
6. Pollin B, Laplante S, Cesaro P, Nguyen-Legros J. (1983) Simultaneous visualization of Nuclear yellow and iron-dextran complex for demonstration of branched neurons by retrograde axonal transport. *J Neurosci Methods*, 8, 205.
7. Arbuthnott GW, Wright AK, Hamilton MH, Brown JR. (1982) Orthograde transport of Nuclear yellow: a problem and its solution. *J Neurosci Methods*, 6, 365.
8. Katan S, Gottschall J, Neuhuber W. (1982) Simultaneous visualization of horseradish peroxidase and Nuclear Yellow in tissue sections for neuronal double labeling. *Neurosci Lett*, 28, 121.
9. Bentivoglio M, Kuypers HG, Catsman-Berrevoets CE. (1980) Retrograde neuronal labeling by means of Bisbenzimidazole and Nuclear Yellow (Hoechst S 769121). Measures to prevent diffusion of the tracers out of retrogradely labeled neurons. *Neurosci Lett*, 18, 19.
10. Bentivoglio M, Kuypers HG, Catsman-Berrevoets CE. (1980) Retrograde neuronal labeling by means of Bisbenzimidazole and Nuclear Yellow (Hoechst S 769121). Measures to prevent diffusion of the tracers out of retrogradely labeled neurons. *Neurosci Lett*, 18, 19.

Disclaimer: This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact our technical service representative for more information.