

Cell Explorer™ Live Cell Labeling Kit

Orange Fluorescence with 405 nm Excitation

Ordering Information	Storage Conditions	Instrument Platform
Product Number: 22616 (2 plates)	Keep in freezer Protect from moisture & light	Fluorescence microscope Flow cytometer

Introduction

Our Cell Explorer™ Live Cell Labeling Kits are a set of tools which can be used to label cells for fluorescence microscopic and flow cytometric investigations of cellular functions. The effective labeling of cells provides a powerful method for studying cellular events in a spatial and temporal context. This particular kit is designed to uniformly label live cells in orange fluorescence. The kit uses our proprietary non-fluorescent CytoCalcein™ Violet 550 dye that becomes strongly fluorescent upon entering into live cells. CytoCalcein™ Violet 550 is a hydrophobic compound that easily permeates intact live cells. The hydrolysis of non-fluorescent CytoCalcein™ Violet 550 by intracellular esterases generates a strongly fluorescent hydrophilic product that is well-retained in the cell cytoplasm. The orange fluorescence dye has the spectral properties of large Stokes Shift at Ex/Em = ~400/550 nm. When well excited with the Violet Laser at 405 nm, the dye emits intense orange fluorescence at ~550 nm. The kit is optimized to be used with a flow cytometer equipped with a Violet Laser and particularly suitable for multicolor flow cytometric analysis of cells. It can also be used with a fluorescence microscope with a customized filter set.

This Cell Explorer™ Live Cell labeling kit provides all the essential components with an optimized cell-labeling protocol (Ex/Em= 405/550 nm). It is an excellent tool to label cells for fluorescence microscopic and flow cytometric investigations of cellular functions. The effective labeling of cells provides a powerful method for studying cellular events in a spatial and temporal context. The kit is useful for a variety of studies, including cell adhesion, chemotaxis, multidrug resistance, cell viability, apoptosis and cytotoxicity. It is suitable for proliferating and non-proliferating cells

Kit Components

Components	Amount
Component A: CytoCalcein™ Violet 550	1 vial
Component B: HHBS (Hanks' buffer with 20 mM Hepes)	1 bottle (100 mL)

Protocol for one 96-well plate

Brief Summary

Prepare cells in growth medium → Remove growth medium → Add CytoCalcein™ Violet 550 working solution 100 µL/well for a 96-well plate or 25 µL/well for a 384-well plate → Stain the cells at 37°C for 30 minutes to 1 hour → Wash and examine the specimen under microscope at Ex/Em= 405/550 nm

Note: Thaw all the components at room temperature before opening.

1. Prepare cells:

- 1.1 For adherent cells: Plate cells overnight in growth medium at 10,000 to 40,000 cells/well/100 µL for a 96-well plate or 2,500 to 10,000 cells/well/25 µL for a 384-well plate.
- 1.2 For non-adherent cells: Centrifuge the cells from the culture medium and then suspend the cell pellets in culture medium at 50,000-100,000 cells/well/100 µL for a 96-well poly-D lysine plate or 10,000-25,000 cells/well/25 µL for a 384-well poly-D lysine plate. Centrifuge the plate at 800 rpm for 2 minutes with brake off prior to the experiment.

Note: Each cell line should be evaluated on an individual basis to determine the optimal cell density.

2. Prepare CytoCalcein™ Violet 550 stain solution:

- 2.1 Prepare CytoCalcein™ Violet 550 stock solution: Add 20µL of DMSO into CytoCalcein™ Violet 550 vial (Component A), and mix them well.

Note: 10 µL of CytoCalcein™ Violet 550 stock solution is enough for 1 plate. Unused CytoCalcein™ Violet 550 stock solution can be aliquoted and stored at < -20 °C for one month if the tubes are sealed tightly. Avoid repeated freeze-thaw cycles and protect it from light.

- 2.2 Prepare CytoCalcein™ Violet 550 working solution: Add 20 µL of DMSO-reconstituted CytoCalcein™ Violet 550 stock solution (from Step 2.1) into 10 mL of HHBS(Component B), and mix them well. The working solution is stable for at least 2 hours at room temperature.

3. Stain the cells:

- 3.1 Remove the growth medium.
3.2 Add 100 µL/well (96-well plate) or 25 µL/well (384-well plate) CytoCalcein™ Violet 550 working solution (from Step 2.2) into the cell plate.
3.3 Incubate the cells in a 37 °C, 5% CO₂ incubator for 30 minutes to 1 hour.
3.4 Remove the CytoCalcein™ Violet 550 working solution from the cells, and wash the cells with HHBS (Component B) for 2 to 3 times, and replace with HHBS.
3.5 Image the cells using a fluorescence microscope with a filter set at Ex/Em = 405/550 nm.

Note: Alternatively, fix the cells at this point. Store the fixed cells at 4 °C and image the cells later.




Figure 1. Image of HeLa cells with Cell Explorer™ Live Cell Labeling Kit in a Costar black wall/clear bottom 96-well plate.

References

1. Wolff M, Wiedenmann J, Nienhaus GU, Valler M, Heilker R. (2006) Novel fluorescent proteins for high-content screening. *Drug Discov Today*, 11, 1054.
2. Lee S, Howell BJ. (2006) High-content screening: emerging hardware and software technologies. *Methods Enzymol*, 414, 468.
3. Haasen D, Schnapp A, Valler MJ, Heilker R. (2006) G protein-coupled receptor internalization assays in the high-content screening format. *Methods Enzymol*, 414, 121.

Warning: This kit is only sold to end users. Neither resale nor transfer to a third party is allowed without written permission from AATBioquest®. Chemical analysis of the kit components is strictly prohibited. Please call us at 408-733-1055 or e-mail us at info@aatbio.com if you have any questions.