

Catalog Number: 100501, 100502, 150704, 150705, 151459, 195109, 195110

Collagenase

Description

Collagenases degrade native helical collagen fibrils. For an extensive review and bibliography see Harris and Krane (1974), Seifter and Harper (1970 and 1971) and Harper (1980).

The substrate, collagen, is the major fibrous component of animal extracellular connective tissue: skin, tendon, blood vessels, bone, etc. Its structural intricacies must be appreciated for an understanding of collagenolysis. (See review by Seifter and Gallop, 1966. See also Robertson and Miller (1972) on cartilagenous collagen.) In brief, collagen consists of fibrils composed of laterally aggregated, polarized tropocollagen molecules (MW 300,000). Each rod-like tropocollagen unit consists of three helically wound polypeptide α -chains around a single axis. The strands have repetitive glycine residues at every third position and an abundance of proline and hydroxyproline. The amino acid sequence is characteristic of the tissue of origin. Tropocollagen units combine uniformly in a lateral arrangement reflecting charged and uncharged amino acids along the molecule thus creating an axially repeating periodicity (See also Ninmi 1968). Cross linkages continue to develop and collagen becomes progressively more insoluble and resistant to lysis on aging (Hamlin et al. 1975; Harris and Farrell 1972). See also Piez and Torchia (1975), Bruns and Gross (1973) and Tanzer (1973).

Gelatin results when soluble tropocollagen is denatured, for example on mild heating, and the polypeptide chains become randomly dispersed. In this state the strands are vulnerable to cleavage by a wide variety of proteases.

True collagenase may cleave simultaneously across all three chains or attack a single strand (Seifter and Harper 1970). The enzyme has an important role in connective tissue metabolism (Dresden 1971) and is produced by specific cells involved in repairs and remodelling processes (Wahl et al. 1975; Robertson et al. 1972; Sakamoto et al. 1975a). Table I in the review by Seifter and Harper (1971) lists known collagenases. Under normal conditions the mammalian enzyme is bound to α 2-macroglobulin or other serum antiproteases (Eisen et al. 1971). Where the enzyme is functional as in the postpartum uterus, it may be obtained directly from the tissue homogenate (Ryan and Woessner 1971). Mammalian collagenases

split collagen in its native triple-helical conformation at a specific site yielding fragments, TCA and TCB representing 3/4 and 1/4 lengths of the tropocollagen molecule. (Woolley et al. 1975b; Gross et al. 1974). After fragmentation the pieces tend to uncoil into random coil polypeptides (gelatin).

The first mammalian enzyme to be studied was that from a culture of tadpole tail tissue. (Nagai et al. 1966; Skai and Gross 1967; Bauer et al. 1971; Harper et al. 1971; Harper and Gross 1972). Collagenases from human and other mammalian sources have been reported on by Bauer et al. (1975), Brady (1975), Huang and Abramson (1975), Sakamoto et al. (1975a and b), Wahl et al. (1985), Wooley et al. (1975), Fujiwara et al. (1974), Gross et al. (1974), Werb and Burleigh (1974), Sakamoto et al. (1973), Woessner and Ryan (1973), Tokoro et al. (1972), Robertson and Miller (1972), Vaes (1972), Bauer et al. (1971), Donoff et al. (1971), Eisen et al. (1971, 1970, and 1968), Ryan and Woessner (1971), Lazarus et al. (1968). Chesney et al. (1974) report on human platelet collagenase. Eisen et al. (1973) report finding a collagenase in the fiddler crab hepato-pancreas and Phillips and Dresden (1973) in a land planarium. There also exist bacterial collagenases, usually from host invasive strains. These enzymes differ from mammalian collagenases in that they attack many sites along the helix (Seifter and Harper 1971). Collagenases from *Clostridium histolyticum*, first prepared by Mandl et al. (1953), have been most thoroughly studied. Welton and Woods (1975 and 1973) and Keil et al. (1975) report on a collagenase from *Achromobacter iophagus*, Carrick and Berk (1975) and Schoellmann and Fisher (1966) on one from *Pseudomonas aeruginosa* and Merkel et al. (1975) on collagenase from a marine bacterium.

Of particular interest is the relationship between active collagenase and the pathology of rheumatoid arthritis. It has been suggested that the normally inhibited collagenase in articular structures may be activated thereby causing the characteristic tissue destruction in joints. (Abe and Nagai 1973; Kruze and Wojtecka 1972; Bauer et al. 1971; Evanson et al. 1968). Rheumatoid synovial fluid contains an activating material not found in joint fluid from patients with osteoarthritis (Harris and Krane 1974). Harris et al. (1969) reported that the collagenase inhibiting capacity of rheumatoid arthritic patients is reduced by one half. They also reported on two synovial collagenases. See also Woolley et al. (1975a and b), Harris and McCroskery (1974) and Leibovich and Weiss (1971).

Collagenases have also been associated with invasive tumors in rabbits. (Harris et al. 1972; McCroskery et al. 1975 and 1973).

Collagenase has found widespread application in the isolation of specific

cell types from attendant connective tissue (Kono 1969; Peck 1964). For example, there is much interest in the experimental transplantation of pancreatic islet cells to alleviate diabetic symptoms (Barker 1975). See also Ewart et al. (1975), Shibato et al. (1975), Gates et al. (1972), Ashcroft et al. (1971) and Lacy and Kostianovsky (1967). Intact parenchymal cells from rat liver tissue have also been isolated. (Seglen 1973 and 1972; Berry and Friend 1969; Howard et al. 1967). Kitabchi et al. (1971) isolated rat adrenal cells and Czech and Fain (1971) fat cells. DeOca and Malinin (1975) report on primary cell cultures from human kidney and Trusler (1975) on a G-banding technique for human chromosomes. See also Benya et al. (1973) for many additional references.

Particular enzymatic activities of crude collagenases have correlated with the tissues from which the cells were obtained (or with the uses to which the cells are put), and as a result of the correlations several formal types have been established.

Type 1

Containing average amounts of assayed activities (collagenase, caseinase, clostripain, and tryptic activities). It is generally recommended for fat, adrenal, and liver cells.

Type 2

Containing greater clostripain activity. It is generally used for heart, bone, muscle, thyroid, cartilage, and liver cells.

Type 3

Containing low proteolytic activity. It is usually used for mammary and fetal cells.

Type 4

Containing low tryptic activity. It is commonly used for islets and other applications where receptor integrity is crucial.

The wide spectrum of activity of crude collagenase makes it especially useful as a debriding agent. Such clinical applications were highlighted at a symposium reported by Mandl (1972). See also Howes et al. (1959) and Hamit and Upjohn (1960).

Sussman (1968) has indicated its possible value for the treatment of

herniated discs. He found that on injection into the intervertebral space a selective dissolution of the nucleus pulposus and fibrocartilaginous tissue is effected.

Characteristics of Collagenase from *Cl. histolyticum*:

Crude preparations contain not only several collagenases but also a sulphydryl protease, clostripain (Mitchell 1968), a trypsin-like enzyme (Peterkofsky and Dregelmann 1971; Sparrow and McQuade 1973) and an aminopeptidase (Kessler and Yaron 1973).

Sugasawara and Harper (1984) and Bond and Van Wart (1984) report on purification of the collagenases of *Cl. histolyticum*.

Molecular weight: Harper et al. (1965), isolated two fractions, A and B. Their molecular weights are 105,000 and 57,400 respectively.

Composition: Mandl et al. (1964) reported two distinct collagenases: one very active on native collagen and synthetic substrates but with negligible activity on gelatin and azocoll, and a second active on gelatin but less active on synthetic substrates. Schaub and Strauch (1968) also purified two collagenases: Clostridiopeptidase A and Collagenase 2, as did Yoshida and Noda (1965); Collagenases I and II. Harper and Kang (1970) and Mitchell (1968b) also report two collagenases. Kono (1968) describes three fractions: Aa and Ba and Bb. It has been hypothesized that the enzymes are composed of subunits: a tetramer of approximately 105,000 M. W. and a dimer of 57,000. The presence of zinc or cobalt in the molecule is also indicated. (Harper et al. 1965; Takahashi and Seifter 1970). Amino acid compositions have been reported by Mandl et al. (1964), Harper et al. (1965) and Yoshida and Noda (1965). The enzymes contain neither cysteine nor methionine (Takahashi and Seifter 1970).

Optimum pH: 7-9 for the A-a enzyme (Kono et al., 1968).

Activators: Ca^{2+} is required. (Takahashi and Seifter, 1970).

Inhibitors: Metal chelating agents such as cysteine, EDTA or o-phenanthroline but not DFP. It is also inhibited by α 2-macroglobulin - a large plasma glycoprotein (Werb et al. 1974). Nagase et al. (1983) and Stricklin and Welgus (1983) report on natural collagenase inhibitors. The human skin enzyme is inhibited by human serum (Eisen et al. 1970) but granulocyte collagenase is not (Lazarus et al. 1968). Human serum contains α 2-macroglobulin and α 1-antitrypsin that may inhibit certain collagenases as well as a third inhibitor reported by Woolley et al. (1975).

It has also been shown that collagenase A is photo-inactivated in the presence of methylene blue (Takahashi and Seifter, 1970) See also Karakiulakis et al., 1991.

Specificity: Clostridial collagenase I or clostridiopeptidase A degrades the helical regions in native collagen preferentially at the Y-Gly bond in the sequence -Pro-Y-Gly-Pro- where Y is most frequently a neutral amino acid. This bond in synthetic peptide substrates may also be split. See also Soberano and Schoellmann (1972a) and Harris and Farrell (1972). The specificities of the three Kono enzymes have been studied (Kono o, 1968).

Applications

For collagen structural and biosynthetic studies researchers generally use more highly purified collagenase preparations free of other proteolytic activities.

Clostridiopeptidase A (MP catalog # 194124), also called collagenase 1, is a highly purified enzyme essentially free of collagenase 2, clostripain, and other proteases.

For tissue dissociation most researchers employ either crude collagenase preparations or chromatographically purified collagenase combined with other enzymes such as elastase, trypsin, and/or papain.

Typical Assay

Assay methods utilizing labelled collagen have been reported by Gisslow and McBride (1975), Robertson et al. (1972) and Sakamoto et al. (1972). Since true collagenase attacks the helical region of the molecule, change in optical rotary dispersion reflects collagen degradation (Keil et al. 1975).

Method: A modification of the procedure of Mandl et al. (1953). Collagenase is incubated for 5 hours with native collagen. The extent of collagen breakdown is determined using the Moore and Stein (1948) colorimetric ninhydrin method. Amino acids liberated are expressed as micromoles leucine per milligram collagenase. One unit equals one micromole of L-leucine equivalents from collagen in 5 hours at 37°C and pH 7.5 under the specified conditions.

Reagents

- 0.05 M TES [tris(hydroxymethyl)-methyl-2-aminoethane sulfonate] buffer with 0.36 mM calcium chloride, pH 7.5
- 4% Ninhydrin in methyl cellosolve
- 0.2 M Sodium citrate with 0.71 mM stannous chloride, pH 5.0
- Ninhydrin-citric acid mixture: Prepare by mixing 50 ml of the 4% ninhydrin in methyl cellosolve with 50 ml of 0.2 M citrate with 0.71 mM stannous chloride, pH 5.0. Allow mixture to stir for 5 minutes.
- 50% n-Propanol
- Substrate: bovine achilles tendon Collagen and vitamin free casein
- 50% (w/v) Trichloroacetic acid

Enzyme

Dissolve enzyme at a concentration of 1 mg/ml in 0.05 M TES with 0.36 mM calcium chloride, pH 7.5. Dilutions run are 1/10 and 1/20 in the above buffer.

Procedure

Weigh 25 mg of bovine collagen into each of four test tubes. Include at least two tubes to serve as blanks which will contain no enzyme. Add 5.0 ml of 0.05 M TES buffer to the tubes and incubate at 37°C for 15 minutes. Start the reaction by adding 0.1 ml of enzyme dilution to appropriate tubes.

After 5 hours, stop the collagenase reaction by transferring 0.2 ml of solution (leaving behind the collagen) to test tubes containing 1.0 ml of ninhydrin-citric acid mixture. Include an enzyme blank (collagen incubated with 0.1 ml TES buffer in place of enzyme). Heat for 20 minutes in a boiling water bath. After cooling, dilute with 5 ml of 50% n-propanol. Let stand for 15 minutes and read absorbance at 600 nm. From an L-leucine standard curve determine micromoles amino acid equivalent to leucine liberated.

Non-specific protease activity (i. e. caseinase activity) is determined using the above assay and substituting 25 milligrams vitamin free casein for collagen. The reaction is stopped after 5 hours by the addition of 0.5 ml of 50% trichloroacetic acid. After centrifugation, 0.2 ml of the supernatant is transferred to 1.0 ml of ninhydrin and treated as above. Caseinase activity is calculated as collagenase activity.

Calculation

Units/mg = (micromoles L-leucine equivalents liberated) / (mg enzyme in

digestion mixture)

Assay for Clostridiopeptidase A

Collagenase I - *Cl. histolyticum*

Method: The assay is based on the enzyme-specific splitting of a Pz-Pro-Leu-Gly-Pro-Arg substrate between leucine and glycine (Wüncch and Heidrich, 1963). This results in a colored lipophilic fragment and a non-colored tripeptide. The change in absorbance of the fragment at 320 nm per unit time is a measure of enzyme activity.

Reagents

- 0.1 M Tris*HCl, pH 7.1
- 0.1 M Calcium chloride
- 25 mM Citric acid, pH 3.5
- Ethyl acetate
- Anhydrous sodium sulfate
- Substrate: Pz-Pro-Leu-Gly-Pro-Arg (molecular weight 812.93). Dissolve to 5 mg/ml in methanol. Dilute to 1 mg/ml with Tris buffer.
- Standard: Pz-Pro-Leu (molecular weight 466.54). Dissolve to 5 mg/ml in methanol. Dilute to 1 mg/ml with Tris buffer.

Enzyme

Stock solution: Dissolve at 1 mg/ml in reagent grade water. Dilute stock 1:20 and 1:50 for assay.

Procedure

	Micrograms of Standard						
	0	50	100	200	400	600	800
Standard	0 ml	0.05 ml	0.1 ml	0.2 ml	0.4 ml	0.6 ml	0.8 ml
Tris buffer	1.1 ml	1.05 ml	1.0 ml	0.9 ml	0.7 ml	0.5 ml	0.3 ml
Calcium chloride	0.2 ml	0.2 ml	0.2 ml	0.2 ml	0.2 ml	0.2 ml	0.2 ml

Mix well and transfer 0.5 ml to labeled tubes containing 1 ml citric acid and 5 ml ethyl acetate. Vortex for 15 seconds. All the phases separate at room temperature. Transfer the organic phase to tubes containing 350

mg sodium sulfate. Shake gently to allow ethyl acetate to dry. Read the A_{320} of the dried ethyl acetate vs. air.

Determine the net A_{320} (test - blank) and plot A_{320} nm/ug standard from the slope of the curve. Determine the calibration factor as follows:

$$\text{Factor} = (1) / (A_{320} \text{ nm/ug} \times 466 \text{ ug/umole})$$

Enzyme Assay: Pipette 0.2 ml calcium chloride and 1.0 ml substrate into a series of numbered tubes. Incubate in a water bath at 25°C to achieve temperature equilibration. At timed intervals, add 0.1 ml enzyme dilution to the respective tubes. Include 2 tubes with 0.1 ml water as blanks. Incubate exactly 15 minutes at 25°C and at timed intervals withdraw 0.5 ml and transfer to tubes containing 1 ml citric acid and 5 ml ethyl acetate. Proceed as with standard curve, reading A_{320} of dried ethyl acetate vs. air. Determine net A_{320} .

Calculation

$$\text{Units/mg} = (\text{Net } A_{320} \times \text{dilution} \times \text{Factor}) / (0.1 \times 15 \times \text{mg/ml in stock})$$

Availability:

Catalog Number	Description	Sizes Available
194124	Clostridiopeptidase A from Clostridium histolyticum. Activity approximately 5 units/dry weight. Supplied as a lyophilized powder tested for BAEE activities and casein digestion.	5 U 10 U
362001	Collagenase from Clostridium histolyticum. Activity approximately 300 Mandl units/mg. This mixture is suited for dissolving connective tissue and exposing embedded cells without destroying the cell membranes and other structures, since the content of unspecific proteases is very low. The high purity and complete absence of pathogenic spores makes this enzyme best suited for cell research and cell culture applications.	300 KU
100501	Collagenase from Clostridium histolyticum. Sterile, salt-free,	50 mg

	lyophilized. Activity approximately 125-250 units/mg dry weight.	
150705	Collagenase from <i>Clostridium</i> <i>histolyticum</i> . Cell Culture Reagent.	1 gm 5 x 1 gm
151459	Collagenase from <i>Clostridium</i> <i>histolyticum</i> , highly purified, free of detectable non-specific proteases. Activity 2,000 - 3,000 units per ml when reconstituted with one ml buffer.	1 vial
150704	Collagenase from <i>Clostridium</i> <i>histolyticum</i> . Type 3, Activity approximately 3500 u/gm; clostripain, trypsin and caseinase \geq 100 u/gm (by casein-folin method, pH 7.0)	1 gm 5 x 1 gm
195109	Collagenase from <i>Clostridium</i> <i>histolyticum</i> , Grade I, salt-free, lyophilized. Activity approximately 125-220 units per mg solid.	25 mg 100 mg 500 mg 1 gm
100502	Collagenase from <i>Clostridium</i> <i>histolyticum</i> , Grade II, lyophilized powder. Activity approximately 125 - 250 units/mg dry weight	25 mg 100 mg 500 mg 1 gm
195110	Collagenase from <i>Clostridium</i> <i>histolyticum</i> , Type 4, salt-free, lyophilized. Activity approximately 125 units per mg solid. Hepatocyte Isolation Grade, contains clostripain, neutral protease and trypsin activities.	25 mg 100 mg 500 mg

References

1. "An Editorial: Collagenase and Rheumatoid Arthritis," *New Eng. J. Med.*, v. **279**, 942 (1968).
2. Abe, S., and Nagai, Y.: "Evidence for the Presence of a Complex of Collagenase with $[[\alpha]]$ 2-Macroglobulin in Human Rheumatoid Synovial Fluid: A Possible Regulatory Mechanism of Collagenase Activity in Vivo," *J. Biochem. Japan*, v. **73**, 897 (1973).

3. Adams, E., Antoine, S., and Golstein, A.: "Collagenase Action on the Synthetic Tripeptide Polymers, (Gly-Pro-Ala)_n and (Gly-Ala-Pro)_n," *Biochim. Biophys. Acta*, v. **185**, 251 (1969).
4. Amsterdam, A., and Jamieson, J.: "Studies on Dispersed Pancreatic Exocrine Cells. I. Dissociation Technique and Morphologic Characteristics of Separated Cells," *J. Cell Biol.*, v. **63**, 1037 (1974).
5. Amsterdam, A., and Jamieson, J.: "Studies on Dispersed Pancreatic Exocrine Cells. II. Functional Characteristics of Separated Cells," *J. Cell Biol.*, v. **63**, 1057 (1974).
6. Ashcroft, S., Bassett, J., and Randle, P.: "Isolation of Human Pancreatic Islets Capable of Releasing Insulin and Metabolising Glucose in Vitro," *Lancet I*, v. **888** (1971).
7. Bagchi, N.: "What Makes Silica Toxic," *Brit. J. Industrial Med.*, v. **49**, 163 (1992).
8. Barker, C.: "Transplantation of the Islets of Langerhans and the Histocompatibility of Endocrine Tissue," *Diabetes*, v. **24**, 766 (1975).
9. Barrett, A., Knight, C., Brown, M., and Tisljar, U.: "A Continuous Fluorimetric Assay for Clostridial Collagenase and Pz-peptidase Activity," *Biochem. J.*, v. **260**, 259 (1989).
10. Bauer, E., Eisen, A., and Jeffrey, J.: "Immunologic Relationship of a Purified Human Skin Collagenase to Other Human and Animal Collagenases," *Biochim. Biophys. Acta*, v. **206**, 152 (1970).
11. Bauer, E., Eisen, A., and Jeffrey, J.: "Studies on Purified Rheumatoid Synovial Collagenase *in Vitro* and *in Vivo*," *J. Clin. Invest.*, v. **50**, 2056 (1971).
12. Bauer, E., Jeffrey, J., and Eisen, A.: "Preparation of Three Vertebrate Collagenases in Pure Form," *Biochem. Biophys. Res. Comm.*, v. **44**, 813 (1971).
13. Bauer, E., Stricklin, G., Jeffrey, J., and Eisen, A.: "Collagenase Production by Human Skin Fibroblasts," *Biochem. Biophys. Res. Comm.*, v. **64**, 232 (1975).
14. Bensusan, H.: "An Investigation of the Products of an Enzymatic Hydrolysis of Collagens," *Biochem.*, v. **8**, 4716 (1969).
15. Benya, P., Berger, K., Golditch, M., and Schneir, M.: "Recommendations Concerning the Use of Bacterial Collagenase for Isolating Collagen-Associated Molecule from Tissue," *Anal. Biochem.*, v. **53**, 313 (1973).
16. Bergmann, U., Michaelis, J., Oberhoff, R., Knauper, V., Beckmann, R., and Tshesche, H.: "Enzyme Linked Immunosorbent Assays (ELISA) for the Quantitative Determination of Human Leukocyte Collagenase and Gelatinase (Original Paper)," *J. Clin. Chem. Clin. Biochem.*, v. **27**, 351 (1989).

17. Berry, L., and Shuttleworth, C.: "Bacterial Collagenase and Collagen Identification," *Connective Tissue Research*, v. 17, 153 (1988).
18. Berry, M., and Friend, D.: "High-Yield Preparation of Isolated Rat-Liver Parenchymal Cells," *J. Cell Biol.*, v. 43, 506 (1969).
19. Bicsak, T., and Harper, E.: "Purification and Characterization of Tadpole Back-skin Collagenase with Low Gelatinase Activity," *J. Biol. Chem.*, v. 259, 13145 (1984).
20. Bienkowski, R., Curran, S., and Berg, R.: "Kinetics of Intracellular Degradation of Newly Synthesized Collagen," *Biochem.*, v. 25, 2455 (1986).
21. Blankaert, A., Mazieres, B., Eeckhout, Y., and Vaes, G.: "Direct Extraction and Assay of Collagenase from Human Osteoarthritic Articular Cartilage," *Clin. Chim. Acta*, v. 185, 73 (1989).
22. Bols, M., Binderup, L., Hansen, J., and Rasmussen, P.: "Inhibition of Collagenase by Aranciamycin and Aranciamycin Derivatives," *J. Med. Chem.*, v. 35, 2768 (1992).
23. Bond, M., and VanWart, H.: "Purification and Separation of Individual Collagenases of *Clostridium histolyticum* Using Red Dye Ligand Chromatography," *Biochem.*, v. 23, 3077 (1984).
24. Bond, M., and VanWart, H.: "Characterization of the Individual Collagenases from *Clostridium histolyticum*," *Biochem.*, v. 23, 3085 (1984).
25. Bond, M., and VanWart, H.: "Relationship between the Individual Collagenases of *Clostridium histolyticum*: Evidence for Evolution by Gene Duplication," *Biochem.*, v. 23, 3092 (1984).
26. Brady, A.: "Collagenase in Scleroderma," *J. Clin. Invest.*, v. 56, 1175 (1975).
27. Bruns, R., and Gross, J.: "Band Pattern of the Segment-Long-Spacing Form of Collagen. Its Use in the Analysis of Primary Structure," *Biochem.*, v. 12, 809 (1973).
28. Callaway, J., Garcia, J., Hersh, C., Yeh, R., and Gilmore-Herbert, M.: "Use of Lectin Affinity Chromatography for the Purification of Collagenase from Human Polymorphonuclear Leukocytes," *Biochem.*, v. 25, 4757 (1986).
29. Carrick, L., and Berk, R.: "Purification and Partial Characterization of a Collagenolytic Enzyme from *Pseudomonas aeruginosa*," *Biochim. Biophys. Acta*, v. 391, 422 (1975).
30. Chesney, C., Harper, E., and Colman, R.: "Human Platelet Collagenase," *J. Clin. Invest.*, v. 53, 1647 (1974).
31. Chu, K.: "Collagenase Chemonucleolysis via Epidural Injection: A Review of 252 Cases," *Clin. Orthop. Related Res.*, v. 215, 99 (1987).
32. Connell, J., and Rousselot, L.: "Use of Enzymatic Agents in Debridement of Burn and Wound Sloughs," *Surgery*, v. 30, 43 (1951).

33. Czech, M., and Fain, J.: "Dactinomycin Inhibition of Dexamethasone Action on Glucose Metabolism in White Fat Cells," *Biochim. Biophys. Acta*, v. **230**, 185 (1971).
34. DeBellis, R., Mandl, I., MacLennan, J., and Howes, E.: "Separation of Proteolytic Enzymes of *Cl. histolyticum*," *Nature*, v. **174**, 1191 (1954).
35. DeOca, H., and Malinin, T.: "Dispersion and Cultivation of Renal Cells After Short-Term Storage of Kidneys," *J. Clin. Microbiol.*, v. **2**, 243 (1975).
36. Dono, K., Gotoh, M., Fukuzaki, T., Ohzato, H., Kanai, T., Monden, M., and Mori, T.: "Low-Temperature Collagenase Digestion - An Improved Islet Isolation Method from Cold Preserved Pancreas," *Transplantation Proc.*, v. **24**, 1511 (1992).
37. Donoff, R., McLenan, J., and Grillo, H.: "Preparation and Properties of Collagenases from Epithelium and Mesenchyme of Healing Mammalian Wounds," *Biochim. Biophys. Acta*, v. **227**, 639 (1971).
38. Dresden, M.: "Evidence for the Role of Collagenase in Collagen Metabolism," *Nature New Biol.*, v. **231**, 55 (1971).
39. Eisen, A., Bauer, E., and Jeffrey, J.: "Human Skin Collagenase. The Role of Serum Alpha-Globulins in the Control of Activity *in Vivo* and *in Vitro*," *Proc. Nat. Acad. Sci. USA* v. **68**, 248 (1971).
40. Eisen, A., Bloch, K., and Sakai, T.: "Inhibition of Human Skin Collagenase by Human Serum," *J. Lab Clin. Med.*, v. **75**, 258 (1970).
41. Eisen, A., Henderson, K., Jeffrey, J., and Bradshaw, R.: "A Collagenolytic Protease from the Hepatopancreas of the Fiddler Crab *Uca pugilator*. Purification and Properties," *Biochem.*, v. **12**, 1814 (1973).
42. Eisen, A., Jeffrey, J., and Gross, J.: "Human Skin Collagenase, Isolation and Mechanism of Attack on the Collagen Molecule," *Biochim. Biophys. Acta*, v. **151**, 637 (1968).
43. Ellis, S., Nabeshima, K., and Biswas, C.: "Monoclonal Antibody Preparation and Purification of a Tumor Cell Collagenase-Stimulatory Factor," *Cancer Res.*, v. **49**, 3385 (1989).
44. Elsback, P., and Rizack, M.: "The Effect of Collagenase Preparations Contaminated with Phospholipase C Activity on Adipose Tissue Lecithin," *Biochim. Biophys. Acta*, v. **198**, 82 (1970).
45. Emöd, I., Tong, N., and Keil, B.: "Chemical Characterization of the Homogenous Collagenase from *Clostridium histolyticum*," *Biochem. Biophys. Acta*, v. **659**, 283 (1981).
46. Evanson, J., Jeffrey, J., and Krane, S.: "Studies on Collagenase from Rheumatoid Synovium in Tissue Culture," *J. Clin. Invest.*, v. **47**, 263 (1968).

47. Ewadh, M., Tudball, N., and Rose, F.: "Homocysteine Uptake by Human Umbilical Vein Endothelial Cells in Culture," *Biochim. Biophys. Acta*, v. **1054**, 263 (1990).
48. Ewart, R., Kornfield, S., and Kipnis, D.: "Effect of Lectins on Hormone Release from Isolated Rat Islets of Langerhans," *Diabetes*, v. **24**, 705 (1975).
49. Foley, J., and Aftonomos, B.: "Growth of Human Breast Neoplasms in Cell Culture," *J. Natl. Cancer Inst.*, v. **34**, 217 (1965).
50. Fox, C., Stanford, J., and Sampath, A.: "Topical Use of Collagenase with Silver Sulfadiazine in Burns," in *Collagenase*, (Mandl, I., ed.), Gordon and Breach, NY, 145 (1972).
51. French, M., Mookhtiar, K., and Van Wart, H.: "Limited Proteolysis of Type I Collagen at Hyperreactive Sites by Class I and II *Clostridium histolyticum* Collagenases: Complementary Digestion Patterns," *Biochem.*, v. **26**, 681 (1987).
52. Fujiwara, K., Sakai, T., Oda, T., and Igarashi, S.: "Demonstration of Collagenase Activity in Rat Liver Homogenate," *Biochem. Biophys. Res. Comm.*, v. **60**, 166 (1974).
53. Galardy, R., and Grobelny, D.: "Inhibition of Collagenase from *Clostridium histolyticum* by Phosphoric and Phosphonic Amides," *Biochem.*, v. **22**, 4556 (1983).
54. Gallop, P., Seifter, S., and Meilman, E.: "Studies on Collagen. I. The Partial Purification, Assay, and Mode of Activation of Bacterial Collagenase," *J. Biol. Chem.*, v. **227**, 891 (1957).
55. Gates, R., Hunt, M., Smith, R., and Lazarus, N.: "Return to Normal of Blood-Glucose, Plasma Insulin, and Weight Gain in New Zealand Obese Mice after Implantation of Islets of Langerhans," *Lancet*, **II**, 567 (1972).
56. Gisslow, M., and McBride, B.: "A Rapid Sensitive Collagenase Assay," *Anal. Biochem.*, v. **68**, 70 (1975).
57. Gottlieb, A., Peterkofsky, B., and Udenfriend, S.: "Use of Collagenase to Identify Enzymatically Formed Collagen and a Hydroxyproline-deficient Intermediate," *J. Biol. Chem.*, v. **240**, 3099 (1965).
58. Grant, N., and Alburn, H.: "Studies on the Collagenase of *Clostridium histolyticum*," *Arch. Biochem. Biophys.*, v. **82**, 245 (1959).
59. Grant, N., and Alburn, H.: "Collagen Solubilization by Mammalian Proteinases," *Arch. Biochem. Biophys.*, v. **89**, 262 (1960).
60. Gray, D., McShane, P., Grant, A., and Morris, R.: "A Method for Isolation of Islets of Langerhans from the Human Pancreas," *Diabetes*, v. **33**, 1055 (1984).

61. Grobelny, D., and Galardy, R. : "Aldehyde and Ketone Substrate Analogues Inhibit the Collagenase of *Clostridium histolyticum*," *Biochemistry*, v. **24**, 6145 (1985).
62. Gross, J., Harper, E., Harris, E., McCroskery, P., Highberger, J., Corbett, C., and Kang, A. : "Animal Collagenases: Specificity of Action, and Structures of the Substrate Cleavage Site," *Biochem. Biophys. Res. Comm.*, v. **61**, 605 (1974).
63. Hamit, H., and Upjohn, H. : "Debridement of Experimental Burns with *Clostridium histolyticum* H4 Enzyme," *Ann. Surg.*, v. **151**, 589 (1960).
64. Hamlin, C., Kohn, R., and Luschin, J. : "Apparent Accelerated Aging of Human Collagen in Diabetes Mellitus," *Diabetes*, v. **24**, 902 (1975).
65. Harper, E. : "Collagenases," *Ann. Rev. Biochem.*, v. **49**, 1063 (1980).
66. Harper, E., and Gross, J. : "Collagenase, Procollagenase and Activator Relationships in Tadpole Tissue Cultures," *Biochem. Biophys. Res. Comm.*, v. **48**, 1147 (1972).
67. Harper, E., and Kang, A. : "Studies on the Specificity of Bacterial Collagenase," *Biochem. Biophys. Res. Comm.*, v. **41**, 482 (1970).
68. Harper, E., Bloch, K., and Gross, J. : "The Zymogen of Tadpole Collagenase," *Biochem.*, v. **10**, 3035 (1971).
69. Harper, E., Seifter, S., and Hospelhorn, V. : "Evidence for Subunits in Bacterial Collagenase," *Biochem. Biophys. Res. Comm.*, v. **18**, 627 (1965).
70. Harris, E., and Farrell, M. : "Resistance to Collagenase: A Characteristic of Collagen Fibrils Cross-Linked by Formaldehyde," *Biochim. Biophys. Acta.*, v. **278**, 133 (1972).
71. Harris, E., and Krane, S. : "Collagenases," *New Eng. J. Med.*, v. **291**, 605 (1974).
72. Harris, E., and McCroskery, P. : "Degradation of Cartilage Collagen by Rheumatoid Synovial Collagenase," *New Eng. J. Med.*, v. **290**, 1 (1974).
73. Harris, E., DiBona, D., and Drane, S. : "Collagenases of Human Synovial Fluid," *J. Clin. Invest.*, v. **48**, 2104 (1969).
74. Harris, E., Faulkner, C., and Wood, S. : "Collagenase in Carcinoma Cells," *Biochem. Biophys. Res. Comm.*, v. **48**, 1247 (1972).
75. Hausmann, E., and Kaufman, E. : "Collagenase Activity in a Particulate Fraction from *Bacteroides melaninogenicus*," *Biochim. Biophys. Acta*, v. **194**, 612 (1969).
76. Hayward, N. : "Collagenase Maceration of Kidney for Micro-dissection," *Nature*, v. **186**, 403 (1960).
77. Heindl, M., Femandjian, S., and Keil, B. : "Circular Dichroism Comparative Studies of Two Bacterial Collagenases and Thermolysin," *Biochim. Biophys. Acta*, v. **624**, 51 (1980).

78. Hemminki, K. : "An Improved Method for Preparing Brain Cell Suspensions," *FEBS Lett.*, v. 9, 290 (1970).
79. Hersh, C., Hey, R., Callaway, J., Garcia, J., and Gilmore-Herbert, M. : "Induction of Collagenase Production in U937 Cells by Phorbol Ester and Partial Purification of the Induced Enzyme," *Biochem.*, v. 25, 4750 (1986).
80. Hilfer, S., and Brown, J. : "Collagenase. Its Effectiveness as a Dispersing Agent for Embryonic Chick Thyroid and Heart," *Exp. Cell Res.*, v. 65, 246 (1970).
81. Hinz, R., and Syverton, J. : "Mammalian Cell Cultures for Study of Influenza Virus. I. Preparation of Monolayer Cultures with Collagenase," *Proc. Soc. Exp. Biol. Med.*, v. 101, 19 (1959).
82. Hoover, N., and Ivins, J. : "Wound Debridement," *Arch. Surg.*, v. 79, 701 (1959).
83. Houck, J., and Patel, Y. : "Collagenolytic Activity of Mammalian Pancreas," *Proc. Soc. Exp. Biol. Med.*, v. 116, 382 (1964).
84. Howard, R., Christensen, A., Gibbs, F., and Pesch, L. : "The Enzymatic Preparation of Isolated Intact Parenchymal Cells from Rat Liver," *J. Cell Biol.*, v. 35, 675 (1967).
85. Howes, E. : "Use of Proteolytic Enzymes as Debriding Agents," *Surgery*, v. 42, 948 (1957).
86. Howes, E., MacLennan, J., Mandl, I., and De Bellis, R. : "Debridement of Burn Slough by Peptidases Recovered from *Clostridium histolyticum*," *Bull. N.Y. Acad. Med.*, v. 31, 413 (1955).
87. Howes, E., Mandl, I., Zaffuto, S., and Ackermann, W. : "The Use of *Clostridium histolyticum* Enzymes in the Treatment of Experimental Third Degree Burns," *Surg. Gyn. Obs.*, v. 109, 177 (1959).
88. Huang, C., and Abramson, M. : "Purification and Characterization of Collagenase from Guinea Pig Skin," *Biochim. Biophys. Acta*, v. 384, 484 (1975).
89. Jeffrey, J., and Gross, J. : "Collagenase from Rat Uterus. Isolation and Partial Characterization," *Biochem.*, v. 9, 268 (1970).
90. Kang, A., Nagai, Y., Piez, K., and Gross, J. : "Studies on the Structure of Collagen Utilizing a Collagenolytic Enzyme from Tadpole," *Biochem.*, v. 5, 509 (1966).
91. Karakiulakis, G., Papadimitriou, E., Missirlis, E., and Maragoudakis, M. : "Effect of Divalent Metal Ions on Collagenase from *Clostridium histolyticum*," *Biochem. Intl.*, v. 24, 397 (1991).
92. Keil, B. : "Some Newly Characterized Collagenases from Procaryotes and Lower Eucaryotes," *Mol. Cell. Biochem.*, v. 23, 87 (1979).
93. Keil, B., Gilles, A., Lecroisey, A., Hurion, N., and Tong, N. : "Specificity of Collagenase from *Achromobacter lophagus*," *FEBS Lett.*, v. 56, 292 (1975).

94. Keller, S., and Mandl, I.: "The Preparation of Purified Collagenase," *Arch. Biochem. Biophys.*, v. **101**, 81 (1963).
95. Kerwar, S., Nolan, J., Ridge, S., Oronsky, A., and Slakey, L.: "Properties of a Collagenase Inhibitor Partially Purified from Cultures of Smooth Muscle Cells," *Biochim. Biophys. Acta*, v. **632**, 183 (1980).
96. Kessler, E., and Yaron, A.: "A Novel Aminopeptidase from *Clostridium histolyticum*," *Biochem. Biophys. Res. Comm.*, v. **50**, 405 (1973).
97. Kitabachi, A., Wilson, D., and Sharma, R.: "Steroidogenesis in Isolated Adrenal Cells of Rat," *Biochem. Biophys. Res. Comm.*, v. **4**, 898 (1971).
98. Kono, T.: "Purification and Partial Characterization of Collagenolytic Enzyme from *Clostridium histolyticum*," *Biochem.*, v. **7**, 1106 (1968).
99. Kono, T.: "Roles of Collagenases and Other Proteolytic Enzymes in the Dispersal of Animal Tissues," *Biochim. Biophys. Acta*, v. **178**, 397 (1969).
100. Kruze, D, and Wojtcka, E.: "Activation of Leucocyte Collagenase Proenzyme by Rheumatoid Synovial Fluid," *Biochim. Biophys. Acta*, v. **258**, 436 (1972).
101. Kula, M., Hatef-Haghi, D., Tauber-Finkelstein, M., and Shaltiel, S.: "Consecutive Use of W-Aminoalkylagaroses. Resolution and Purification of Clostripain and Collagenase from *Clostridium histolyticum*," *Biochem. Biophys. Res. Comm.*, v. **69**, 389 (1976).
102. Lacy, P.E., and Kostianovsky, M.: "Method for the Isolation of Intact Islets of Langerhans from the Rat Pancreas," *Diabetes*, v. **16**, 35 (1967).
103. Lasfargues, E.: "Cultivation and Behavior *in Vitro* of the Normal Mammary Epithelium of the Adult Mouse," *Anat. Rec.*, v. **127**, 117 (1957).
104. Lasfargues, E., and Ozzello, L.: "Cultivation of Human Breast Carcinomas," *J. Natl. Cancer Inst.*, v. **21**, 1131 (1958).
105. Lasfarques, E.: "Collagenase-Treated Skin Autografts for Rapid Healing of Burns," in *Collagenase*, (Mandl, I., ed)., Gordon and Breach, NY, 151 (1972).
106. Lazarus, G., Brown, R., Daniels, J., and Fullmer, H.: "Human Granulocyte Collagenase," *Science*, v. **159**, 1483 (1968).
107. Lazarus, G., Daniels, J., Brown, R., Bladen, H., and Fullmer, H.: "Degradation of Collagen by a Human Granulocyte Collagenolytic System," *J. Clin. Invest.*, v. **47**, 2622 (1968).
108. Lee-Own, V., and Anderson, J.: "The Preparation of a Bacterial Collagenase Containing Negligible Non-Specific Protease Activity," *Prep. Biochem.*, v. **5**, 229 (1975).

109. Lefebvre, V., and Vaes, G. : "The Enzymatic Evaluation of Procollagenase and Collagenase Inhibitors in Crude Biological Media," *Biochim. Biophys. Acta*, v. **992**, 355 (1989).
110. Leibovich, S., and Weiss, J. : "Failure of Human Rheumatoid Synovial Collagenase to Degrade Either Normal or Rheumatoid Arthritic Polymeric Collagen," *Biochim. Biophys. Acta*, v. **251**, 109 (1971).
111. Lin, M., and Chen, Y. : "Effect of Copper Ion on Collagenase Release - Its Implication in Corneal Vascularization," *Inv. Ophthalm. Visual Sci.*, v. **33**, 558 (1992).
112. Liotta, L., Lanzer, W., and Garbisa, S. : "Identification of a Type V Collagenolytic Enzyme," *Biochem. Biophys. Res. Comm.*, v. **98**, 184 (1981).
113. Lwebuga-Mukasa, J., Harper, E., and Taylor, P. : "Collagenase Enzymes from *Clostridium*: Characterization of Individual Enzymes," *Biochem. USA*, v. **15**, 4736 (1976).
114. MacLennan, J., Mandl, I., and Howes, E. : "Bacterial Digestion of Collagen," *J. Clin. Invest.*, v. **32**, 1317 (1953).
115. Makinen, P., and Makinen, K. : "Near Stoichiometric, Irreversible Inactivation of Bacterial Collagenases by o-Chloranil (3,4,5,6-Tetrachloro-1,2-benzoquinone)," *Biochim. Biophys. Res. Comm.*, v. **153**, 74 (1988).
116. Mallya, S., Mookhtiar, K., and VanWart, H. : "Kinetics of Hydrolysis of Type-I, Type-II, and Type-III Collagens by the Class-I and Class-II *Clostridium histolyticum* Collagenases," *J. Protein Chem.*, v. **11**, 99 (1992).
117. Mandl, I. : "Collagenases and Elastases," in *Advances in Enzymology*, **23**, (Nord, F., ed.), Interscience Publ., NY, 163 (1961).
118. Mandl, I. : "Collagenase," *Science*, v. **169**, 1234 (1970).
119. Mandl, I., ed. : *Collagenase*, published by Gordon and Breach, NY, (1972).
120. Mandl, I., Keller, S., and Manahan J. : "Multiplicity of *Clostridium histolyticum* Collagenases," *Biochem.*, v. **3**, 1737 (1964).
121. Mandl, I., MacLennan, J., Howes, E., DeBellis, R., and Sohler, A. : "Isolation and Characterization of Proteinase and Collagenase from *Cl. histolyticum*," *J. Clin. Invest.*, v. **32**, 1323 (1953).
122. Mandl, I., Zipper, H., and Ferguson, L. : "Clostridium histolyticum Collagenase: Its Purification and Properties," *Arch. Biochem. Biophys.*, v. **74**, 465 (1958).
123. McCroskery, P., Richards, J., and Harris, E. : "Purification and Characterization of Collagenase Extracted from Rabbit Tumours," *Biochem. J.*, v. **152**, 131 (1975).

124. McCroskery, P., Wood, S., and Harris, E. : "Gelatin: A Poor Substrate for Mammalian Collagenase," *Science*, v. **182**, 70 (1973).
125. Merkel, J., Dreisbach, J., and Ziegler, H. : "Collagenolytic Activity of Some Marine Bacteria," *Appl. Microbiol.*, v. **29**, 145 (1975).
126. Michaels, S., Gallop, P., Seifter, S., and Meilman, E. : "Specificity of Collagenase," *Biochim. Biophys. Acta*, v. **29**, 450 (1958).
127. Mitchell, W. : "Pseudocollagenase: A Protease from *Clostridium histolyticum*," *Biochim. Biophys. Acta*, v. **159**, 554 (1968).
128. Miyahara, M., Hayashi, K., Berger, J., Tanzawa, K., Njieha, F.K., Treislard, R.L., and Prockop, D.J. : "Formation of Collagen Fibrils by Enzymatic Cleavage of Precursors of Type I Collagen *in Vitro*," *J. Biol. Chem.*, v. **259**, 9891 (1984).
129. Mooktiar, K., Steinbrink, D., and VanWart, H. : "Mode of Hydrolysis of Collagen-like Peptides by Class I and Class II *Clostridium histolyticum* Collagenases: Evidence for both Endopeptidase and Tripeptidyl-carboxypeptidase Activities," *Biochem.*, v. **24**, 6527 (1985).
130. Moore, S., and Stein, W. : "Photometric Ninhydrin Method," *J. Biol. Chem.*, v. **176**, 367 (1948).
131. Moore, W., and Spilburg, C. : "Purification of Human Collagenases with a Hydroxamic Acid Affinity Column," *Biochem.*, v. **25**, 5189 (1986).
132. Morris, G. : "A High Molecular Weight Collagenase Inhibitor Made by Rabbit Chondrocytes in Cell Culture," *Matrix*, v. **9**, 127 (1989).
133. Murphy, G., McAlpine, C., Poll, C., and Reynolds, J. : "Purification and Characterization of a Bone Metalloproteinase that Degrades Gelatin and Types IV and V Collagen," *Biochim. Biophys. Acta*, v. **831**, 49 (1985).
134. Nagai, Y. : "Specificity of Collagenase. The Action of Collagenase on Synthetic Peptides," *J. Biochem.*, v. **50**, 493 (1961).
135. Nagai, Y., Lapiere, C., and Gross, J. : "Tadpole Collagenase: Preparation and Purification," *Biochem.*, v. **5**, 3123 (1966).
136. Nagase, H., and Harris, E. : "Ovostatin: A Novel Proteinase Inhibitor From Chicken Egg White. II. Mechanisms of Inhibition Studied with Collagenase and Thermolysin," *J. Biol. Chem.*, v. **258**, 7490 (1983).
137. Nagase, H., Cawston, T., DeSilva, M., and Barret, A. : "Identification of Plasma Kallikrein as an Activator of Latent Collagenase in Rheumatoid Synovial Fluid," *Biochim. Biophys. Acta*, v. **702**, 133 (1982).

138. Nagase, H., Harris, E., Woessner, J., and Brew, K. : "Ovostatin: A Novel Proteinase Inhibitor from Chicken Egg White. I. Purification, Physicochemical Properties and Tissue Distribution of Ovostatin," *J. Biol. Chem.*, v. **258**, 7481 (1983).
139. Nagase, H., Jackson, R., Brinckerhoff, C., Vater, C., and Harris, E. : "A Precursor Form of Latent Collagenase Produced in a Cell-free System with mRNA from Rabbit Synovial Cells," *J. Biol. Chem.*, v. **256**, 11951 (1981).
140. Narinesingh, D., and Ngo, T. : "Enzyme Amplified Potentiometric Assay for Collagenase Using Urease-Labelled, Solid Supported Substrate," *Anal. Letters*, v. **22**, 323 (1989).
141. Nedergaard, J., and Lindberg, O. : "The Brown Fat Cell," *Intl. Rev. Cytology*, v. **74**, 187 (1982).
142. Niedermuller, H., and Petersen, R. : "Modification of Collagenase Activity in the Skin of Young, Adult and Old Rats by a Plasma Factor," *Arch. Gerontology and Geriatrics*, **Suppl. 3**, 277 (1992).
143. Nimni, M. : "A Defect in the Intramolecular and Intermolecular Cross-linking of Collagen Caused by Penicillamine," *J. Biol. Chem.*, v. **243**, 1457 (1968).
144. Nixon, J., Bottomley K., Broadhurst, M., Brown, P., Johnson, W., Lawton, G., Marley, J., Sedgwick, A., and Wilkinson, S. : "Potent Collagenase Inhibitors Prevent Interleukin-1-Induced Cartilage Degradation *In vitro*," *Intl. J. Tissue Reactions-Exp. Clin. Aspects*, v. **13**, 237 (1991).
145. Noel, J., Rabinovitch, A., Olson, L., Kyriakides, G., Miller, J., and Mintz, D. : "A Method for Large-Scale, High-Yield Isolation of Canine Pancreatic Islets of Langerhans," *Metabolism*, v. **31**, 184 (1982).
146. Peck, W., Birge, S., and Fedak, S. : "Bone Cells: Biochemical and Biological Studies After Enzymatic Isolation," *Science*, v. **146**, 1476 (1964).
147. Peterkofsky, B., and Diegelmann, R. : "Use of a Mixture of Proteinase-Free Collagenases for the Specific Assay of Radioactive Collagen in the Presence of Other Proteins," *Biochem.*, v. **10**, 988 (1971).
148. Phillips, J., and Dresden, M. : "A Collagenase in Extracts of the Invertebrate *Bipalium kewense*," *Biochem. J.*, v. **133**, 329 (1973).
149. Piez, K., and Torchia, D. : "Possible Contribution of Ionic Clustering to Molecular Packing of Collagen," *Nature*, v. **258**, 87 (1975).

150. Pollard, M., and Stan, T. : "Propagation of Duck Hepatitis in Tissue Culture Prepared with Collagenase," *J. Bacteriol.*, v. **80**, 571 (1960).
151. Robertson, P., and Miller, E. : "Cartilage Collagen: Inability to Serve as a Substrate for Collagenases Active Against Skin and Bone Collagen," *Biochim. Biophys. Acta*, v. **289**, 247 (1972).
152. Robertson, P., Taylor, R., and Fullmer, H. : "A Reproducible Quantitative Collagenase Radiofibril Assay," *Clin. Chim. Acta*, v. **42**, 43 (1972).
153. Rodbell, M. : "Metabolism of Isolated Fat Cells. I. Effects of Hormones on Glucose Metabolism and Lipolysis," *Journ. Biol. Chem.*, v. **239**, 375 (1964).
154. Ryan, J., and Woessner, F. : "Mammalian Collagenase: Direct Demonstration in Homogenates of Involuting Rat Uterus," *Biochem. Biophys. Res. Comm.*, v. **44**, 144 (1971).
155. Sakai, T., and Cross, J. : "Some Properties of the Products of Reaction of Tadpole Collagenase with Collagen," *Biochem.*, v. **6**, 518 (1967).
156. Sakamoto, S., Goldhaber, P., and Glimcher, M. : "A New Method for the Assay of Tissue Collagenase," *Proc. Soc. Exp. Biol. Med.*, v. **139**, 1057 (1972).
157. Sakamoto, S., Sakamoto, M., Goldhaber, P., and Glimcher, M. : "Isolation of Tissue Collagenase From Homogenates of Embryonic Chick Bones," *Biochem. Biophys. Res. Comm.*, v. **53**, 1102 (1973).
158. Sakamoto, S., Sakamoto, M., Goldhaber, P., and Glimcher, M. : "Collagenase and Bone Resorption: Isolation of Collagenase from Culture Medium Containing Serum After Stimulation of Bone Resorption by Addition of Parathyroid Hormone Extract," *Biochem. Biophys. Res. Comm.*, v. **63**, 172 (1975a).
159. Sakamoto, S., Sakamoto, M., Goldhaber, P., and Glimcher, M. : "Studies on the Interaction Between Heparin and Mouse Bone Collagenase," *Biochim. Biophys. Acta*, v. **385**, 41 (1975b).
160. Schaub, M., and Strauch, L. : "Evidence for a New Bacterial Collagenolytic Enzyme," *Biochem. Biophys. Res. Comm.*, v. **21**, 34 (1965).
161. Schaub, M., and Strauch, L. : "Purification of Collagenase IV," *Hoppe-Seyler's Z. Physiol. Chem.*, v. **349**, 809 (1968).
162. Schoellman, G., and Fisher, E. : "A Collagenase From *Pseudomonas aeruginosa*," *Biochim. Biophys. Acta*, v. **122**, 557 (1966).
163. Seglen, P. : "Preparation of Rat Liver Cells II. Effects of Ions and Chelators on Tissue Dispersion," *Exp. Cell Res.*, v. **76**, 25 (1973).

164. Seifter, S., and Gallop, P. : in *The Proteins*, Vol. IV, 2nd Ed. , (Neurath, H., ed.), 238 (1966).
165. Seifter, S., and Harper, E. : "Collagenases," in *Methods in Enzymology XIX* (Perlmann, G. and Lorand, L., eds.), Academic Press, NY, 613 (1970).
166. Seifter, S., and Harper, E. : "Collagenases," in *The Enzymes III*, (Boyer, P.D., ed.), Academic Press, NY, 649 (1971).
167. Seifter, S., Gallop, P., Klein, L., and Meilman, E. : "Studies on Collagen II. Properties of Purified Collagenase and its Inhibitor," *J. Biol. Chem.*, v. **234**, 285 (1959).
168. Seifter, S., Takahasi, S., and Harper, E. : "Further Demonstration that Cysteine Reacts with the Metal Component of Collagenase," *Biochim. Biophys. Acta*, v. **214**, 559 (1970).
169. Sgaragli, G., Sen, I., Baba, A., Schulz, R., and Cooper, J. : "The Mechanism of Action of Collagenase on the Inhibition of Release of Acetylcholine from Synaptosomal Preparations," *Brain Res.*, v. **134**, 113 (1977).
170. Shibata, A., Ludvigsen, C., and Mcdaniel, M. : "Standardization of a Digestion-Filtration Method for Isolation of Pancreatic Islets," *Diabetes*, v. **24 Supp. 2** (Abstracts), 444 (1975).
171. Soberano, M., and Schoellmann, G. : "Specificity of Bacterial Collagenase: Studies with Peptides Newly Synthesized Using the Solid-Phase Method," *Biochim. Biophys. Acta*, v. **271**, 133 (1972a).
172. Soberano, M., and Schoellmann, G. : "Anomalous Kinetics Observed in the Catalysis of a Peptide Substrate with Collagenase," *Biochim. Biophys. Acta*, v. **289**, 401 (1972b).
173. Sparrow, L., and McQuade, A. : "Isolation by Affinity Chromatography of Neutral Proteinase from *Clostridium histolyticum*," *Biochim. Biophys. Acta*, v. **302**, 90 (1973).
174. Spina, M., Garbin, G., Field, J., and Serafini-Fracassini, A. : "The Application of Collagenase, Purified by Affinity Chromatography, to the Isolation of Insoluble Elastin from Bovine Aorta," *Biochim. Biophys. Acta*, v. **400**, 162 (1975).
175. Stack, M., and Gray, R. : "Comparison of Vertebrate Collagenase and Gelatinase Using a New Fluorogenic Substrate Peptide," *J. Biol. Chem.*, v. **264**, 4277 (1989).
176. Steinbrink, D., Bond, M., and Van Wart, H. : "Substrate Specificity of β -Collagenase from *Clostridium histolyticum*," *J. Biol. Chem.*, v. **260**, 2771 (1985).
177. Steven, F., Knott, J., Jackson, D., and Podrazky, V. : "Collagen-Protein-Polysaccharide Interactions in Human Intervertebral Disc," *Biochim. Biophys. Acta*, v. **188**, 307 (1969).

178. Stricklin, G., and Welgus, H. : "Human Skin Fibroblast Collagenase Inhibitor. Purification and Biochemical Characterization," *J. Biol. Chem.*, v. **258**, 12252 (1983).
179. Sugasawara, R., and Harper, E. : "Purification and Characterization of Three Forms of Collagenase from *Clostridium histolyticum*," *Biochem.*, v. **23**, 5175 (1984).
180. Sussman, B. : "Intervertebral Discolysis with Collagenase," *J. Nat. Med. Assoc.*, v. **60**, 184 (1968).
181. Takahashi, S., and Seifter, S. : "Dye-Sensitized Photo-Inactivation of Collagenase A," *Biochim. Biophys. Acta*, v. **214**, 556 (1970).
182. Tanaka, T., Metori, K., Mineo, S., Hirotsu, M., Furuya, T., Matsumoto, H., Satoh, T., and Kobayashi, S. : "Studies on Collagenase Inhibitors. 4. Inhibitors of Bacterial Collagenase in *Coptidis rhizoma* - Note," *J. Pharm. Soc. Japan*, v. **111**, 538 (1991).
183. Tanzer, M. : "Cross-Linking of Collagen," *Science*, v. **180**, 561 (1973).
184. Taylor, A., Levy, B., and Simpson, J. : "Collagenolytic Activity of Sarcoma Tissues in Culture," *Nature*, v. **228**, 366 (1970).
185. Thompson, M., Shuplock, N., and Fetterman, S. : "Collagenase Method of Renal Microdissection," *Arch. Pathol.*, v. **78**, 568 (1964).
186. Tokoro, Y., Eisen, A., and Jeffrey, J. : "Characterization of a Collagenase from Rat Skin," *Biochim. Biophys. Acta*, v. **258**, 289 (1972).
187. Tong, N., Tsugita, A., and Keil-Dlouha, V. : "Purification and Characterization of Two High-molecular-mass Forms of *Achromobacter* Collagenase," *Biochim. Biophys. Acta*, v. **874**, 296 (1986).
188. Treadwell, B., Neidel, J., Pavia, M., Towle, C., Trice, M., and Mankin, H. : "Purification and Characterization of Collagenase Activator Protein Synthesized by Articular Cartilage," *Arch. Biochem. Biophys.*, v. **251**, 715 (1986).
189. Treadwell, B., Towle, C., Ishizue, K., Mankin, K., Pavia, M., Ollivierre, F., and Gray, D. : "Stimulation of the Synthesis of Collagenase Activator Protein in Cartilage by a Factor Present in Synovial-Conditioned Medium," *Arch. Biochem. Biophys.*, v. **251**, 724 (1986).
190. Trusler, S. : "G Bands Produced by Collagenase," *Lancet*, I, 44 (1975).
191. Vaes, G. : "The Release of Collagenase as an Inactive Proenzyme by Bone Explants in Culture," *Biochem. J.*, v. **126**, 275 (1972).
192. Vaitkevicius, V., Simpson, W., and Brennan, M. : "Freeze Preservation of Enzymatically Prepared Tumor Cell Suspensions and Their Use in Tumor Transplantation," *Cancer Res.*, v. **21**, 187 (1961).

193. VanWart, H., and Steinbrink, D. : "A Continuous Spectrophotometric Assay for *Clostridium histolyticum* Collagenase," *Anal. Biochem.*, v. **113**, 356 (1981).
194. Van Wart, H., and Steinbrink, D. : "Complementary Substrate Specificities of Class I and Class II Collagenases from *Clostridium histolyticum*," *Biochem.*, v. **24**, 6520 (1985).
195. Vencill, C., Rasnick, D., Crumley, K., Nishino, N., and Powers, J. : "*Clostridium histolyticum* Collagenase: Development of New Thio Ester, Fluorogenic, and Depsipeptide Substrates and New Inhibitors," *Biochem.*, v. **24**, 3149 (1985).
196. von Hippel, P., Gallop, P., Seifter, S., and Cunningham, R. : "An Enzymatic Examination of the Structure of the Collagen Macromolecule," *J. Am. Chem. Soc.*, v. **82**, 2774 (1960).
197. Wahl, L., Wahl, S., Mergenhagen, S., and Martin, G. : "Collagenase Production by Lymphokine-Activated Macrophages," *Science*, v. **187**, 261 (1975).
198. Warren, G., and Gray, J. : "A Simplified Culture Medium for the Production of Collagenase," *Nature*, v. **192**, 755 (1961).
199. Weingarten, H., Marten, R., and Feder, J. : "Synthetic Substrates of Vertebrate Collagenase," *Biochem.*, v. **24**, 6730 (1985).
200. Welgus, H., and Stricklin, G. : "Human Skin Fibroblast Collagenase Inhibitor. Comparative Studies in Human Connective Tissues, Serum and Amniotic Fluid," *J. Biol. Chem.*, v. **258**, 12259 (1983).
201. Welgus, H., Jeffrey, J., and Eisen, A. : "The Collagen Substrate Specificity of Human Fibroblast Collagenase," *J. Biol. Chem.*, v. **256**, 9516 (1981).
202. Welton, R., and Woods, D. : "Halotolerant Collagenolytic Activity of *Achromobacter lophagus*," *J. Gen. Microbiol.*, v. **75**, 191 (1973).
203. Welton, R., and Woods, D. : "Collagenase Production by *Achromobacter lophagus*," *Biochim. Biophys. Acta*, v. **384**, 228 (1975).
204. Werb, Z., and Burleigh, M. : "A Specific Collagenase from Rabbit Fibroblasts in Monolayer Culture," *Biochem. J.*, v. **137**, 373 (1974).
205. Werb, Z., Burleigh, M., Barrett, A., and Starkey, P. : "The Interaction of α 2-Macroglobulin with Proteinases," *Biochem. J.*, v. **139**, 359 (1974).
206. Wize, J. : "A Latent Collagenase from Rheumatoid Synovial Fluid. Purification and Partial Characterization," *Biochim. Biophys. Acta*, v. **615**, 199 (1980).

207. Woessner, J., and Ryan, J. : "Collagenase Activity in Homogenates of the Involuting Rat Uterus," *Biochim. Biophys. Acta*, v. **309**, 397 (1973).
208. Wolf, G. : "The Molecular Basis of the Inhibition of Collagenase by Vitamin A," *Nutrition Reviews*, v. **50**, 292 (1992).
209. Wolters, G., Vosscheperkeuter, G., Vandeiijnen, J., and Vanschilfgaarde, R. : "An Analysis of the Role of Collagenase and Protease in the Enzymatic Dissociation of the Rat Pancreas for Islet Isolation," *Diabetologia*, v. **35**, 735 (1992).
210. Woolley, D., Lindberg, K., Glanville, R., and Evanson, J. : "Action of Rheumatoid Synovial Collagenase on Cartilage Collagen," *Eur. J. Biochem.*, v. **50**, 437 (1975b).
211. Woolley, D., Roberts, D., and Evanson, J. : "Inhibition of Human Collagenase Activity by a Small Molecular Weight Serum Protein," *Biochem. Biophys. Res. Comm.*, v. **66**, 747 (1975).
212. Woolley, D., Roberts, D., and Evanson, J. : "Small Molecular Weight b1 Serum Protein which Specifically Inhibits Human Collagenases," *Nature*, v. **261**, 325 (1976).
213. Woolley, F., Glanville, R., Crossley, M., and Evanson, J. : "Purification of Rheumatoid Synovial Collagenase and Its Action on Soluble and Insoluble Collagen," *Eur. J. Biochem.*, v. **54**, 611 (1975a).
214. Wünsch, E., and Heidrich, H. : *Hoppe-Seyler's Z. Physiol. Chem.*, v. **332**, 300 (1963).
215. Wünsch, E., and Heidrich, H. : *Hoppe-Seyler's Z. Physiol. Chem.*, v. **332**, 149 (1963).
216. Yagisawa, S. Morita, F., Nagai, Y., Noda, H., and Ogura, Y. : "Kinetic Studies on the Action of Collagenase," *J. Biochem. Japan*, v. **58**, 407 (1965).
217. Yiotakis, A., Hatgiyannacou, A., Dive, V., and Toma, F. : "New Thiol Inhibitors of *Clostridium histolyticum* Collagenase - Importance of the P3' Position," *Eur. Journ. Biochem.*, v. **172**, 761 (1988).
218. Yoshida, E., and Noda, H. : "Isolation and Characterization of Collagenases I and II from *Clostridium histolyticum*," *Biochim. Biophys. Acta*, v. **105**, 562 (1965).
219. Zimmerman, W. : "The Importance of Collagenase for the Local Treatment of Major Burns," in *Collagenase*, (Mandl, I., ed.), Gordon and Breach, NY, 131 (1972).