


Catalog Number: 102918, 190522, 194831, 198957, 811030, 811032, 811033, 811034, 811036

## Sodium dodecyl sulfate

**Structure:**



**Molecular Formula:** C<sub>12</sub>H<sub>25</sub>NaSO<sub>4</sub>

**Molecular Weight:** 288.38

**CAS #:** 151-21-3

**Synonyms:** SDS; Lauryl sulfate sodium salt; Dodecyl sulfate sodium salt; Dodecyl sodium sulfate; Sodium lauryl sulfate; Sulfuric acid monododecyl ester sodium salt

**Physical Appearance:** White granular powder

**Critical Micelle Concentration (CMC):** 8.27 mM (Detergents with high CMC values are generally easy to remove by dilution; detergents with low CMC values are advantageous for separations on the basis of molecular weight. As a general rule, detergents should be used at their CMC and at a detergent-to-protein weight ratio of approximately ten.<sup>13,14</sup>

**Aggregation Number:** 62

**Solubility:** Soluble in water (200 mg/ml – clear, faint yellow solution), and ethanol (0.1g/10 ml)

**Description:** An anionic detergent<sup>3</sup> typically used to solubilize<sup>8</sup> and denature proteins for electrophoresis.<sup>4,5</sup> SDS has also been used in large-scale phenol extraction of RNA to promote the dissociation of protein from nucleic acids when extracting from biological material.<sup>12</sup> Most proteins bind SDS in a ratio of 1.4 grams SDS to 1 gram protein. The charges intrinsic to the protein become insignificant compared to the overall negative charge provided by the bound SDS. The charge to mass ratio is essentially the same for each protein and will migrate in the gel based only on protein size.

Typical Working Concentration: > 10 mg SDS/mg protein

Typical Buffer Compositions:

*SDS Electrophoresis Gel Running Buffer*

| Component | Composition (g/L) | Molarity or % |
|-----------|-------------------|---------------|
| Tris      | 3.035             | 0.025 M       |
| Glycine   | 14.1              | 0.192 M       |
| SDS       | 1                 | 0.1%          |
| pH 8.3    |                   |               |

*SDS Electrophoresis Gel Sample Solubilization Buffer*

| Component        | Composition (g/L) | Molarity or % |
|------------------|-------------------|---------------|
| Tris             | 15.16             | 0.125 M       |
| SDS              | 20.0              | 2.0%          |
| Sucrose          | 100               | 10.0%         |
| Bromophenol Blue | 0.2               | 0.02%         |

Titrated to pH 8.0 with HCl

Availability:

| Catalog Number | Description   | Size                                    |
|----------------|---|---|
| 102918         | Sodium Dodecyl Sulfate, purity approximately 99%  | 10 g<br>25 g<br>100 g<br>500 g<br>1 kg  |
| 190522         | Sodium Dodecyl Sulfate, purity approximately 95%  | 100 g<br>500 g<br>1 kg                  |
| 194831         | Sodium Dodecyl Sulfate, molecular biology reagent, purity approximately 99%                         | 25 g<br>100 g<br>250 g<br>500 g<br>1 kg |
| 198957         | Sodium Dodecyl Sulfate, 20% Solution; purity approximately 99.5% alkyl sulfate, 97% dodecyl sulfate | 100 ml<br>500 ml                        |

|        |  |       |
|--------|--|-------|
| 811033 | Sodium Dodecyl Sulfate, Ultra Pure, purity | 25 g  |
| 811036 | not less than 99%                          | 50 g  |
| 811034 |  | 100 g |
| 811032 |  | 500 g |
| 811030 |  | 1 kg  |

### References:

1. *Merck Index*, **12th Ed.**, No. 8782.
2. *CRC Handbook of Electrophoresis*, Lewis, L.A. and Oppelt, J.J. (eds.), v. 1, CRC Press: Boca Raton, Florida (1980).
3. *Colloid Polym. Sci.*, v. **272**, 141 (1994).
4. *Electrophoresis* (Weinheim, Fed. Repub. Ger.), v. **14**, 1038 (1993).
5. *Anal. Biochem.*, v. **214**, 490 (1993).
6. Grohs, R.A., et al., *Anal. Chem.*, v. **63**, 384 (1991).
7. Hames, B.D. and Rickwood, D., *Gel Electrophoresis of Proteins: A Practical Approach*, **2nd Ed.**, IRL Press: New York, NY, pp. 8 (1990).
8. Harewood, K. and Wolff, J.S., *Anal. Biochem.*, v. **55**, 573 (1973).
9. Jagus, R., *Meths. Enzymol.*, v. **152**, 296 (1987).
10. Laemmli, U.K., "Cleavage of structural proteins during the assembly of the head of bacteriophage T4." *Nature*, v. **227**, 680-685 (1970).
11. Nelsen, T.B. and Reynolds, J.A., *Meths. Enzymol.*, v. **48**, 3 (1978).
12. Wallace, D.M., *Meths. Enzymol.*, v. **125**, 39 (1987).
13. Womack, M.D. et al., *Biochim. Biophys. Acta*, **733**, 210 (1983)
14. Matson, R.S. and Goheen, S.C., *LC-GC*, **4(7)**, 624 (1986)