

Sulfolobus DNA Polymerase IV



1-800-632-7799
info@neb.com
www.neb.com



M0327S 001120614061

M0327S



100 units **2,000 U/ml** **Lot: 0011206**

RECOMBINANT **Store at -20°C** **Exp: 6/14**

Description: *Sulfolobus* DNA Polymerase IV is a thermostable Y-family **lesion-bypass** DNA Polymerase that efficiently synthesizes DNA across a variety of DNA template lesions.

Source: An *E. coli* strain that carries the gene encoding DNA polymerase IV from *Sulfolobus islandicus*.

Applications:

- Synthesis of DNA through DNA lesions (1,2)
- Repair of DNA (1)

Supplied in: 10 mM Tris-HCl (pH 7.4), 100 mM KCl, 1 mM DTT, 0.1 mM EDTA and 50% glycerol.

Reagents Supplied with Enzyme:
10X ThermoPol™ Reaction Buffer.

Reaction Conditions: 1X ThermoPol Reaction Buffer. Incubate at 55°C.

1X ThermoPol Reaction Buffer:

20 mM Tris-HCl
10 mM (NH₄)₂SO₄
10 mM KCl
2 mM MgSO₄
0.1% Triton® X-100
pH 8.8 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTPs into acid insoluble material in 30 minutes at 55°C.

Unit Assay Conditions: 1X ThermoPol Reaction Buffer, 200 μM dNTPs including [³H]-dTTP and 15 nM primed M13mp18.

Supplied in: 10 mM Tris-HCl (pH 7.4), 100 mM KCl, 1 mM DTT, 0.1 mM EDTA and 50% glycerol.

Reagents Supplied with Enzyme:
10X ThermoPol™ Reaction Buffer.

Reaction Conditions: 1X ThermoPol Reaction Buffer. Incubate at 55°C.

1X ThermoPol Reaction Buffer:

20 mM Tris-HCl
10 mM (NH₄)₂SO₄
10 mM KCl
2 mM MgSO₄
0.1% Triton® X-100
pH 8.8 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTPs into acid insoluble material in 30 minutes at 55°C.

Unit Assay Conditions: 1X ThermoPol Reaction Buffer, 200 μM dNTPs including [³H]-dTTP and 15 nM primed M13mp18.

Quality Control Assays

Exonuclease Activity: Incubation of a 50 μl reaction in ThermoPol Reaction Buffer containing a minimum of 20 units of *Sulfolobus* DNA Polymerase IV and 1 μg of a mixture of single and double-stranded [³H] *E. coli* DNA for 4 hours at either 37°C or 55°C releases < 0.1% of the total radioactivity.

Endonuclease Activity: Incubation of a 50 μl reaction in ThermoPol Reaction Buffer containing a minimum of 20 units of *Sulfolobus* DNA Polymerase IV with 1 μg of supercoiled φX174 DNA for 4 hours at either 37°C or 55°C results in < 10% conversion to the nicked form as determined by agarose gel electrophoresis.

Enzyme Properties

3' → 5' Exonuclease: No
5' → 3' Exonuclease: No
Strand displacement: No

Heat inactivation: No

Molecular Weight: 40,274 daltons (theoretical)

Quality Control Assays

Exonuclease Activity: Incubation of a 50 μl reaction in ThermoPol Reaction Buffer containing a minimum of 20 units of *Sulfolobus* DNA Polymerase IV and 1 μg of a mixture of single and double-stranded [³H] *E. coli* DNA for 4 hours at either 37°C or 55°C releases < 0.1% of the total radioactivity.

Endonuclease Activity: Incubation of a 50 μl reaction in ThermoPol Reaction Buffer containing a minimum of 20 units of *Sulfolobus* DNA Polymerase IV with 1 μg of supercoiled φX174 DNA for 4 hours at either 37°C or 55°C results in < 10% conversion to the nicked form as determined by agarose gel electrophoresis.

Enzyme Properties

3' → 5' Exonuclease: No
5' → 3' Exonuclease: No
Strand displacement: No

Heat inactivation: No

Molecular Weight: 40,274 daltons (theoretical)

Activity in NEBuffers:

NEBuffer 1	25%
NEBuffer 2	100%
NEBuffer 3	100%
NEBuffer 4	100%

Notes:

Activity at Different Temperatures:

30°C	<5%
37°C	20%
45°C	40%
55°C	95%
65°C	90%
72°C	70%

Half-life @ 95°C: 6 minutes

References:

1. Boudsocq, F. et al. (2001) *Nucleic Acids Res.* 29, 4607-4616.
2. McDonald, JP. et al. (2006) *Nucleic Acids Res.* 34, 1102-1111.

(see other side)

CERTIFICATE OF ANALYSIS

Sulfolobus DNA Polymerase IV



1-800-632-7799
info@neb.com
www.neb.com



M0327S 001120614061

M0327S



100 units **2,000 U/ml** **Lot: 0011206**

RECOMBINANT **Store at -20°C** **Exp: 6/14**

Description: *Sulfolobus* DNA Polymerase IV is a thermostable Y-family **lesion-bypass** DNA Polymerase that efficiently synthesizes DNA across a variety of DNA template lesions.

Source: An *E. coli* strain that carries the gene encoding DNA polymerase IV from *Sulfolobus islandicus*.

Applications:

- Synthesis of DNA through DNA lesions (1,2)
- Repair of DNA (1)

Activity in NEBuffers:

NEBuffer 1	25%
NEBuffer 2	100%
NEBuffer 3	100%
NEBuffer 4	100%

Notes:

Activity at Different Temperatures:

30°C	<5%
37°C	20%
45°C	40%
55°C	95%
65°C	90%
72°C	70%

Half-life @ 95°C: 6 minutes

References:

1. Boudsocq, F. et al. (2001) *Nucleic Acids Res.* 29, 4607-4616.
2. McDonald, JP. et al. (2006) *Nucleic Acids Res.* 34, 1102-1111.

(see other side)

CERTIFICATE OF ANALYSIS

Companion Products Sold Separately:

Magnesium Sulfate (MgSO₄) Solution
#B1003S 6.0 ml

Diluent E
#B8005S 4.0 ml

ThermoPol Reaction Buffer Pack
#B9004S 6.0 ml

ThermoPol II (Mg-free) Reaction Buffer Pack
#B9005S 6.0 ml

ThermoPol DF (Detergent-free) Reaction Buffer Pack
#B9013S 6.0 ml

Deoxynucleotide Solution Set
#N0446S 25 µmol each

Deoxynucleotide Solution Mix
#N0447S 8 µmol each
#N0447L 40 µmol each

THERMOPOL™ is a trademark of New England Biolabs, Inc.

TRITON® is a registered trademark of Union Carbide Corporation.



Companion Products Sold Separately:

Magnesium Sulfate (MgSO₄) Solution
#B1003S 6.0 ml

Diluent E
#B8005S 4.0 ml

ThermoPol Reaction Buffer Pack
#B9004S 6.0 ml

ThermoPol II (Mg-free) Reaction Buffer Pack
#B9005S 6.0 ml

ThermoPol DF (Detergent-free) Reaction Buffer Pack
#B9013S 6.0 ml

Deoxynucleotide Solution Set
#N0446S 25 µmol each

Deoxynucleotide Solution Mix
#N0447S 8 µmol each
#N0447L 40 µmol each

THERMOPOL™ is a trademark of New England Biolabs, Inc.

TRITON® is a registered trademark of Union Carbide Corporation.

