

pBR322 DNA



1-800-632-7799
info@neb.com
www.neb.com



N3033S 091120214021

N3033S

50 µg Lot: 0911202 Exp: 2/14

1,000 µg/ml Store at -20°C

Description: pBR322 DNA is a commonly used plasmid cloning vector in *E. coli* (1). The molecule is a double-stranded circle 4,361* base pairs in length (2). pBR322 contains the genes for resistance to ampicillin and tetracycline, and can be amplified with chloramphenicol. The molecular weight is 2.83 x 10⁶ daltons.

Supplied in: 10 mM Tris-HCl (pH 8.0),
1 mM EDTA.

Preparation: pBR322 is isolated from *E. coli* ER2420 (dam⁺dcm⁺ EcoK M⁻) by a standard plasmid purification procedure.

References:

1. Bolivar, F., Rodriguez, R. L., Greene, P. J., Betlach, M. C., Heynecker, H.L. and Boyer, H.W. (1977) *Gene* 2, 95–113.
2. Watson, N. (1988) *Gene* 70, 399–403

*Sequencing data from Watson (confirmed at New England Biolabs, Inc.) has shown pBR322 to be 4,361 base pairs, not 4,363 base pairs as previously reported.

CERTIFICATE OF ANALYSIS

pBR322 DNA



1-800-632-7799
info@neb.com
www.neb.com



N3033S 091120214021

N3033S

50 µg Lot: 0911202 Exp: 2/14

1,000 µg/ml Store at -20°C

Description: pBR322 DNA is a commonly used plasmid cloning vector in *E. coli* (1). The molecule is a double-stranded circle 4,361* base pairs in length (2). pBR322 contains the genes for resistance to ampicillin and tetracycline, and can be amplified with chloramphenicol. The molecular weight is 2.83 x 10⁶ daltons.

Supplied in: 10 mM Tris-HCl (pH 8.0),
1 mM EDTA.

Preparation: pBR322 is isolated from *E. coli* ER2420 (dam⁺dcm⁺ EcoK M⁻) by a standard plasmid purification procedure.

References:

1. Bolivar, F., Rodriguez, R. L., Greene, P. J., Betlach, M. C., Heynecker, H.L. and Boyer, H.W. (1977) *Gene* 2, 95–113.
2. Watson, N. (1988) *Gene* 70, 399–403

*Sequencing data from Watson (confirmed at New England Biolabs, Inc.) has shown pBR322 to be 4,361 base pairs, not 4,363 base pairs as previously reported.

CERTIFICATE OF ANALYSIS