### Sodium Fluoride (Fluoride)







## P0759S

1 ml 500 mM Lot: 0051204

Exp: 4/15

Store at -20°C

**Description:** Sodium Fluoride (Fluoride, NaF) is a commonly used general inhibitor for protein phosphoseryl and phosphothreonyl phosphatases (PSPs). Routinely Fluoride, combined with Sodium Orthovanadate (NEB# P0758), a general inhibitor for protein phosphotyrosyl phosphatases (PTPs), is used to preserve the protein phosphorylation state in cells, cell lysates, and protein kinase (PK) assays. These inhibitors should ideally be present in all buffers used in the purification of PKs (1, 2).

Supplied in: Sterile purified water.

Molecular Weight: 41.99 daltons

Purity: >99% pure

Suggested Working Concentration: 5-50 mM

#### References:

- Hardie, D.G. et al. (1993). In D. G. Hardie (Ed.), Protein Phosphorylation: A Practical Approach (pp.62). New York: IRL Press.
- 2. Gordon, J.A. (1991) *Methods in Enzymology* 201. 477-482.

CERTIFICATE OF ANALYSIS

### Sodium Fluoride (Fluoride)



1-800-632-7799 info@neb.com www.neb.com



# P0759S

1 ml 500 mM Lot: 0051204

Exp: 4/15

DO mM Store at –20°C

**Description:** Sodium Fluoride (Fluoride, NaF) is a commonly used general inhibitor for protein phosphoseryl and phosphothreonyl phosphatases (PSPs). Routinely Fluoride, combined with Sodium Orthovanadate (NEB# P0758), a general inhibitor for protein phosphotyrosyl phosphatases (PTPs), is used to preserve the protein phosphorylation state in cells, cell lysates, and protein kinase (PK) assays. These inhibitors should ideally be present in all buffers used in the purification of PKs (1, 2).

Supplied in: Sterile purified water.

Molecular Weight: 41.99 daltons

Purity: >99% pure

Suggested Working Concentration: 5-50 mM

#### References:

- Hardie, D.G. et al. (1993). In D. G. Hardie (Ed.), Protein Phosphorylation: A Practical Approach (pp.62). New York: IRL Press.
- 2. Gordon, J.A. (1991) *Methods in Enzymology* 201, 477-482.

CERTIFICATE OF ANALYSIS