

# MspI



1-800-632-7799  
info@neb.com  
www.neb.com



R0106S 053120514051

## R0106S



5,000 units 20,000 U/ml Lot: 0531205

RECOMBINANT Store at -20°C Exp: 5/14

### Recognition Site:

5' . . . C<sup>▼</sup>C G G . . . 3'  
3' . . . G G C<sup>▲</sup>C . . . 5'

**Source:** An *E. coli* strain that carries the cloned MspI gene from *Moraxella* species (ATCC 49670)

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol.

### Reagents Supplied with Enzyme:

10X NEBuffer 4.

### Reaction Conditions:

1X NEBuffer 4.  
Incubate at 37°C.

### 1X NEBuffer 4:

50 mM potassium acetate  
20 mM Tris-acetate  
10 mM magnesium acetate  
1 mM dithiothreitol  
pH 7.9 @ 25°C

**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

**Diluent Compatibility:** Diluent Buffer A  
50 mM KCl, 10 mM Tris-HCl, 0.1 mM EDTA,  
1 mM dithiothreitol, 200 µg/ml BSA and  
50% glycerol (pH 7.4 @ 25°C).

### Quality Control Assays

**Ligation:** After 100-fold overdigestion with MspI, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

**16-Hour Incubation:** A 50 µl reaction containing 1 µg of DNA and 500 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

**Exonuclease Activity:** Incubation of 300 units of enzyme with 1 µg sonicated <sup>3</sup>H DNA (10<sup>5</sup> cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.15% radioactivity.

### Enzyme Properties

#### Activity in NEBuffers:

NEBuffer 1	75%
NEBuffer 2	100%
NEBuffer 3	50%
NEBuffer 4	100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

**Survival in a Reaction:** A minimum of 0.50 unit is required to digest 1 µg of substrate DNA in 16 hours.

### Heat Inactivation: No

**Note:** MspI is an isoschizomer of HpaII.

Not sensitive to *dam*, *dcm* or mammalian CpG methylation.

When the external C in the sequence CCGG is methylated, MspI and HpaII cannot cleave. However, unlike HpaII, MspI can cleave the sequence when the internal C residue is methylated.

= Time-Saver™ Qualified (See www.neb.com for details).

CERTIFICATE OF ANALYSIS

# MspI



1-800-632-7799  
info@neb.com  
www.neb.com



R0106S 053120514051

## R0106S



5,000 units 20,000 U/ml Lot: 0531205

RECOMBINANT Store at -20°C Exp: 5/14

### Recognition Site:

5' . . . C<sup>▼</sup>C G G . . . 3'  
3' . . . G G C<sup>▲</sup>C . . . 5'

**Source:** An *E. coli* strain that carries the cloned MspI gene from *Moraxella* species (ATCC 49670)

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol.

### Reagents Supplied with Enzyme:

10X NEBuffer 4.

### Reaction Conditions:

1X NEBuffer 4.  
Incubate at 37°C.

### 1X NEBuffer 4:

50 mM potassium acetate  
20 mM Tris-acetate  
10 mM magnesium acetate  
1 mM dithiothreitol  
pH 7.9 @ 25°C

**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

**Diluent Compatibility:** Diluent Buffer A  
50 mM KCl, 10 mM Tris-HCl, 0.1 mM EDTA,  
1 mM dithiothreitol, 200 µg/ml BSA and  
50% glycerol (pH 7.4 @ 25°C).

### Quality Control Assays

**Ligation:** After 100-fold overdigestion with MspI, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

**16-Hour Incubation:** A 50 µl reaction containing 1 µg of DNA and 500 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

**Exonuclease Activity:** Incubation of 300 units of enzyme with 1 µg sonicated <sup>3</sup>H DNA (10<sup>5</sup> cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.15% radioactivity.

### Enzyme Properties

#### Activity in NEBuffers:

NEBuffer 1	75%
NEBuffer 2	100%
NEBuffer 3	50%
NEBuffer 4	100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

**Survival in a Reaction:** A minimum of 0.50 unit is required to digest 1 µg of substrate DNA in 16 hours.

### Heat Inactivation: No

**Note:** MspI is an isoschizomer of HpaII.

Not sensitive to *dam*, *dcm* or mammalian CpG methylation.

When the external C in the sequence CCGG is methylated, MspI and HpaII cannot cleave. However, unlike HpaII, MspI can cleave the sequence when the internal C residue is methylated.

= Time-Saver™ Qualified (See www.neb.com for details).

CERTIFICATE OF ANALYSIS