

SphI



1-800-632-7799
info@neb.com
www.neb.com



R0182S 048120714071

R0182S



500 units 10,000 U/ml Lot: 0481207
RECOMBINANT Store at -20°C Exp: 7/14

Recognition Site:

5'...GCATG[▼]C...3'
3'...C[▲]G[▲]TACG...5'

Source: An *E. coli* strain that carries the cloned SphI gene from *Streptomyces phaeochromogenes* (NRRL B-3559)

Also Available In
High Fidelity (HF™) Format

SphI



1-800-632-7799
info@neb.com
www.neb.com



R0182S 048120714071

R0182S



500 units 10,000 U/ml Lot: 0481207
RECOMBINANT Store at -20°C Exp: 7/14

Recognition Site:

5'...GCATG[▼]C...3'
3'...C[▲]G[▲]TACG...5'

Source: An *E. coli* strain that carries the cloned SphI gene from *Streptomyces phaeochromogenes* (NRRL B-3559)

Also Available In
High Fidelity (HF™) Format

Supplied in: 100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 100 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:
10X NEBuffer 2.

Reaction Conditions: 1X NEBuffer 2.
Incubate at 37°C.

1X NEBuffer 2:
50 mM NaCl
10 mM Tris-HCl
10 mM MgCl₂
1 mM DTT
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility: Diluent Buffer B
300 mM NaCl, 10 mM Tris-HCl, 0.1 mM EDTA, 1 mM DTT, 500 µg/ml BSA and 50% glycerol. (pH 7.4 @ 25°C).

Supplied in: 100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 100 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:
10X NEBuffer 2.

Reaction Conditions: 1X NEBuffer 2.
Incubate at 37°C.

1X NEBuffer 2:
50 mM NaCl
10 mM Tris-HCl
10 mM MgCl₂
1 mM DTT
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility: Diluent Buffer B
300 mM NaCl, 10 mM Tris-HCl, 0.1 mM EDTA, 1 mM DTT, 500 µg/ml BSA and 50% glycerol. (pH 7.4 @ 25°C).

Quality Control Assays

Ligation: After 20-fold overdigestion with SphI, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 25 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 7,500 units of enzyme with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.1% radioactivity

Endonuclease Activity: Incubation of 45 units of enzyme with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reaction buffer resulted in < 20% conversion to RF II.

Blue/White Screening Assay: This enzyme has been tested to determine the integrity of the DNA ends produced after digestion with an excess of enzyme. An appropriate vector is digested at

Quality Control Assays

Ligation: After 20-fold overdigestion with SphI, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 25 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 7,500 units of enzyme with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.1% radioactivity

Endonuclease Activity: Incubation of 45 units of enzyme with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reaction buffer resulted in < 20% conversion to RF II.

Blue/White Screening Assay: This enzyme has been tested to determine the integrity of the DNA ends produced after digestion with an excess of enzyme. An appropriate vector is digested at

a unique site within *lacZ^c* gene with a 10-fold excess of enzyme, ligated, transformed and plated on XGal/IPTG/Amp plates. Successful expression of β-galactosidase is a function of how intact its gene remains after cloning, an intact gene gives rise to a blue colony, an interrupted gene (i.e. degraded DNA end) gives rise to a white colony. Enzymes must produce fewer than 3% white colonies to be Blue/White Certified.

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1	100%
NEBuffer 2	100%
NEBuffer 3	50%
NEBuffer 4	100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

(See other side)

CERTIFICATE OF ANALYSIS

a unique site within *lacZ^c* gene with a 10-fold excess of enzyme, ligated, transformed and plated on XGal/IPTG/Amp plates. Successful expression of β-galactosidase is a function of how intact its gene remains after cloning, an intact gene gives rise to a blue colony, an interrupted gene (i.e. degraded DNA end) gives rise to a white colony. Enzymes must produce fewer than 3% white colonies to be Blue/White Certified.

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1	100%
NEBuffer 2	100%
NEBuffer 3	50%
NEBuffer 4	100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

(See other side)

CERTIFICATE OF ANALYSIS

Survival in a Reaction: A minimum of 0.13 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 70 units of enzyme were inactivated by incubation at 65°C for 20 minutes.

Plasmid Cleavage: Number of units required to cleave 1 µg of supercoiled plasmid DNA in one hour: pBR322 = 2 units pUC 19 = 3 units.


Note: Cleaves to leave a 3' CATG extension which can be efficiently ligated to DNA fragments generated by NlaIII.

Not sensitive to *dam*, *dcm* or mammalian CpG methylation.

Companion Products:

SphI-HF™

#R3182S	500 Units
#R3182L	2,500 Units
#R3182M	2,500 Units

 = Time-Saver™ Qualified (See www.neb.com for details).

U.S. Patent No. 5,262,318

Page 2 (R0182)

Survival in a Reaction: A minimum of 0.13 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 70 units of enzyme were inactivated by incubation at 65°C for 20 minutes.

Plasmid Cleavage: Number of units required to cleave 1 µg of supercoiled plasmid DNA in one hour: pBR322 = 2 units pUC 19 = 3 units.


Note: Cleaves to leave a 3' CATG extension which can be efficiently ligated to DNA fragments generated by NlaIII.

Not sensitive to *dam*, *dcm* or mammalian CpG methylation.

Companion Products:

SphI-HF™

#R3182S	500 Units
#R3182L	2,500 Units
#R3182M	2,500 Units

 = Time-Saver™ Qualified (See www.neb.com for details).

U.S. Patent No. 5,262,318

Page 2 (R0182)