

MluI



1-800-632-7799
info@neb.com
www.neb.com



R0198S 027120814081



R0198S



1,000 units 10,000 U/ml Lot: 0271208

RECOMBINANT Store at -20°C Exp: 8/14

Recognition Site:

5'..A[▼]C G C G T..3'
3'..T G C G C A[▲]..5'

Source: An *E. coli* strain that carries the cloned MluI gene from *Micrococcus luteus* (IFO 12992)

Supplied in: 100 mM NaCl, 10 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X NEBuffer 3.

Reaction Conditions:

1X NEBuffer 3.
Incubate at 37°C.

1X NEBuffer 3:

100 mM NaCl
50 mM Tris-HCl
10 mM MgCl₂
1 mM dithiothreitol
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility: Diluent Buffer A
50 mM KCl, 10 mM Tris-HCl, 0.1 mM EDTA,
1 mM dithiothreitol, 200 µg/ml BSA and
50% glycerol (pH 7.4 @ 25°C)

Reagents Supplied with Enzyme:

10X NEBuffer 3.

Reaction Conditions:

1X NEBuffer 3.
Incubate at 37°C.

1X NEBuffer 3:

100 mM NaCl
50 mM Tris-HCl
10 mM MgCl₂
1 mM dithiothreitol
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility: Diluent Buffer A
50 mM KCl, 10 mM Tris-HCl, 0.1 mM EDTA,
1 mM dithiothreitol, 200 µg/ml BSA and
50% glycerol (pH 7.4 @ 25°C)

Quality Control Assays

Ligation: After 10-fold overdigestion with MluI > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 300 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 100 units of enzyme with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.1% radioactivity.

Endonuclease Activity: Incubation of 40 units of enzyme with 1 µg pUC19 plasmid DNA for 4 hours at 37°C in 50 µl reaction buffer resulted in < 10% conversion to RF II.

Quality Control Assays

Ligation: After 10-fold overdigestion with MluI > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 300 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 100 units of enzyme with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.1% radioactivity.

Endonuclease Activity: Incubation of 40 units of enzyme with 1 µg pUC19 plasmid DNA for 4 hours at 37°C in 50 µl reaction buffer resulted in < 10% conversion to RF II.

Enzyme Properties

Activity in NEBuffers

NEBuffer 1	25%
NEBuffer 2	75%
NEBuffer 3	100%
NEBuffer 4	50%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 0.13 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 65°C for 20 minutes.

Notes: Cleavage of mammalian genomic DNA is blocked by CpG methylation.

= Time-Saver™ Qualified (See www.neb.com for details).

CERTIFICATE OF ANALYSIS

MluI



1-800-632-7799
info@neb.com
www.neb.com



R0198S 027120814081



R0198S



1,000 units 10,000 U/ml Lot: 0271208

RECOMBINANT Store at -20°C Exp: 8/14

Recognition Site:

5'..A[▼]C G C G T..3'
3'..T G C G C A[▲]..5'

Source: An *E. coli* strain that carries the cloned MluI gene from *Micrococcus luteus* (IFO 12992)

Supplied in: 100 mM NaCl, 10 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol.

Enzyme Properties

Activity in NEBuffers

NEBuffer 1	25%
NEBuffer 2	75%
NEBuffer 3	100%
NEBuffer 4	50%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 0.13 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 65°C for 20 minutes.

Notes: Cleavage of mammalian genomic DNA is blocked by CpG methylation.

= Time-Saver™ Qualified (See www.neb.com for details).

CERTIFICATE OF ANALYSIS