

MslI



1-800-632-7799
info@neb.com
www.neb.com



R0571S 008121014101

R0571S



500 units 5,000 U/ml Lot: 0081210

RECOMBINANT Store at -20°C Exp: 10/14

Recognition Site:

5'...CAYNN[▼]NNRTG...3'
3'...GTRNN[▲]NNYAC...5'

Single Letter Code: R = A or G, Y = C or T

Source: An *E. coli* strain that carries the cloned MslI gene from *Moraxella osloensis* (B Qiang)

More Units, New Reaction Buffer

MslI



1-800-632-7799
info@neb.com
www.neb.com



R0571S 008121014101

R0571S



500 units 5,000 U/ml Lot: 0081210

RECOMBINANT Store at -20°C Exp: 10/14

Recognition Site:

5'...CAYNN[▼]NNRTG...3'
3'...GTRNN[▲]NNYAC...5'

Single Letter Code: R = A or G, Y = C or T

Source: An *E. coli* strain that carries the cloned MslI gene from *Moraxella osloensis* (B Qiang)

More Units, New Reaction Buffer

Supplied in: 100 mM KCl, 10 mM Tris-HCl (pH 7.6), 0.1 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:
10X NEBuffer 4.

Reaction Conditions: 1X NEBuffer 4.
Incubate at 37°C.

1X NEBuffer 4:
50 mM potassium acetate
20 mM Tris-acetate
10 mM magnesium acetate
1 mM DTT
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility: Diluent Buffer A
50 mM KCl, 10 mM Tris-HCl, 0.1 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol (pH 7.4 @ 25°C).

Supplied in: 100 mM KCl, 10 mM Tris-HCl (pH 7.6), 0.1 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:
10X NEBuffer 4.

Reaction Conditions: 1X NEBuffer 4.
Incubate at 37°C.

1X NEBuffer 4:
50 mM potassium acetate
20 mM Tris-acetate
10 mM magnesium acetate
1 mM DTT
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility: Diluent Buffer A
50 mM KCl, 10 mM Tris-HCl, 0.1 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol (pH 7.4 @ 25°C).

Quality Control Assays

Ligation: After 50-fold overdigestion with MslI, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 250 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 250 units of enzyme with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.1% radioactivity.

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1 100%
NEBuffer 2 100%
NEBuffer 3 25%
NEBuffer 4 100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Quality Control Assays

Ligation: After 50-fold overdigestion with MslI, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 250 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 250 units of enzyme with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.1% radioactivity.

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1 100%
NEBuffer 2 100%
NEBuffer 3 25%
NEBuffer 4 100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 1.0 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 30 units of enzyme were inactivated by incubation at 65°C for 20 minutes.

Note: Not sensitive to *dam*, *dcm* or mammalian CpG methylation.

= Time-Saver™ Qualified (See www.neb.com for details).

CERTIFICATE OF ANALYSIS

Survival in a Reaction: A minimum of 1.0 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 30 units of enzyme were inactivated by incubation at 65°C for 20 minutes.

Note: Not sensitive to *dam*, *dcm* or mammalian CpG methylation.

= Time-Saver™ Qualified (See www.neb.com for details).

CERTIFICATE OF ANALYSIS