

BspHI



1-800-632-7799
info@neb.com
www.neb.com



R0517S 016120614061

R0517S



500 units 10,000 U/ml Lot: 0161206
RECOMBINANT Store at -20°C Exp: 6/14

Recognition Site:

5'... T[▼]CATGA... 3'
3'... AGTACT[▲]... 5'

Source: An *E. coli* strain that carries the cloned BspHI gene from *Bacillus* species H (D. Hall)

Supplied in: 100 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:
10X NEBuffer 4.

Reaction Conditions: 1X NEBuffer 4.
Incubate at 37°C.

1X NEBuffer 4:
50 mM potassium acetate
20 mM Tris-acetate
10 mM magnesium acetate
1 mM DTT
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility: Diluent Buffer A
50 mM KCl, 10 mM Tris-HCl, 0.1 mM EDTA, 1 mM DTT, 200 µg/ml BSA and 50% glycerol (pH 7.4 @ 25°C)

Quality Controls Assays

Ligation: After 10-fold overdigestion with BspHI, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 50 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction produced in 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 100 units for 4 hours at 37°C in 50 µl assay buffer with 1 µg sonicated ³H DNA (10⁵ cpm/µg) released < 0.1% radioactivity.

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1 10%
NEBuffer 2 100%
NEBuffer 3 50%
NEBuffer 4 **100%**

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

CERTIFICATE OF ANALYSIS

BspHI



1-800-632-7799
info@neb.com
www.neb.com



R0517S 016120614061

R0517S



500 units 10,000 U/ml Lot: 0161206
RECOMBINANT Store at -20°C Exp: 6/14

Recognition Site:

5'... T[▼]CATGA... 3'
3'... AGTACT[▲]... 5'

Source: An *E. coli* strain that carries the cloned BspHI gene from *Bacillus* species H (D. Hall)

Supplied in: 100 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:
10X NEBuffer 4.

Reaction Conditions: 1X NEBuffer 4.
Incubate at 37°C.

1X NEBuffer 4:
50 mM potassium acetate
20 mM Tris-acetate
10 mM magnesium acetate
1 mM DTT
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility: Diluent Buffer A
50 mM KCl, 10 mM Tris-HCl, 0.1 mM EDTA, 1 mM DTT, 200 µg/ml BSA and 50% glycerol (pH 7.4 @ 25°C)

Quality Controls Assays

Ligation: After 10-fold overdigestion with BspHI, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 50 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction produced in 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 100 units for 4 hours at 37°C in 50 µl assay buffer with 1 µg sonicated ³H DNA (10⁵ cpm/µg) released < 0.1% radioactivity.

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1 10%
NEBuffer 2 100%
NEBuffer 3 50%
NEBuffer 4 **100%**

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 0.13 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 50 units of enzyme were incubated at 65°C for 20 minutes.

Note: Blocked by overlapping *dam* methylation.

Companion Products:

dam⁻/dcm⁻ Competent *E. coli*
#C2925H 20 transformation reactions
#C2925I 24 transformation reactions

= Time-Saver™ Qualified (See www.neb.com for details).

Survival in a Reaction: A minimum of 0.13 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 50 units of enzyme were incubated at 65°C for 20 minutes.

Note: Blocked by overlapping *dam* methylation.

Companion Products:

dam⁻/dcm⁻ Competent *E. coli*
#C2925H 20 transformation reactions
#C2925I 24 transformation reactions

= Time-Saver™ Qualified (See www.neb.com for details).

CERTIFICATE OF ANALYSIS