

BcoDI



1-800-632-7799
info@neb.com
www.neb.com



R0542S 00112114111

R0542S



1,000 units **10,000 U/ml** **Lot: 0011211**
RECOMBINANT **Store at -20°C** **Exp: 11/14**

Recognition Site:

5'...GTCTC(N)₁▼...3'
3'...CAGAG(N)₅▲...5'

Source: An *E. coli* strain that carries the cloned BcoDI gene from *Bacteroides coprocola* DSM 17136.

Supplied in: 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X NEBuffer 4

Reaction Conditions:

1X NEBuffer 4.

Incubate at 37°C.

1X NEBuffer 4:

50 mM potassium acetate
20 mM Tris-acetate
10 mM magnesium acetate
1 mM dithiothreitol
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility: Diluent Buffer B
300 mM NaCl, 10 mM Tris-HCl (pH 7.4),
0.1 mM EDTA, 1 mM DTT, 500 µg/ml BSA and
50% glycerol.

Heat Inactivation: 65°C for 20 minutes.

Quality Control Assays

Ligation: After 10-fold overdigestion with BcoDI, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 30 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 200 units of enzyme with 1 µg sonicated [³H] DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.1% radioactivity.

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1 50%
NEBuffer 2 75%
NEBuffer 3 75%
NEBuffer 4 **100%**

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 0.13 unit is required to digest 1 µg of substrate DNA in 16 hours.

Notes: BcoDI is an isoschizomer of BsmAI.

Cleavage of mammalian genomic DNA is blocked by some combinations of overlapping CpG methylation.

= Time-Saver™ Qualified (See www.neb.com for details).

CERTIFICATE OF ANALYSIS

BcoDI



1-800-632-7799
info@neb.com
www.neb.com



R0542S 00112114111

R0542S



1,000 units **10,000 U/ml** **Lot: 0011211**
RECOMBINANT **Store at -20°C** **Exp: 11/14**

Recognition Site:

5'...GTCTC(N)₁▼...3'
3'...CAGAG(N)₅▲...5'

Source: An *E. coli* strain that carries the cloned BcoDI gene from *Bacteroides coprocola* DSM 17136.

Supplied in: 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X NEBuffer 4

Reaction Conditions:

1X NEBuffer 4.

Incubate at 37°C.

1X NEBuffer 4:

50 mM potassium acetate
20 mM Tris-acetate
10 mM magnesium acetate
1 mM dithiothreitol
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility: Diluent Buffer B
300 mM NaCl, 10 mM Tris-HCl (pH 7.4),
0.1 mM EDTA, 1 mM DTT, 500 µg/ml BSA and
50% glycerol.

Heat Inactivation: 65°C for 20 minutes.

Quality Control Assays

Ligation: After 10-fold overdigestion with BcoDI, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 30 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 200 units of enzyme with 1 µg sonicated [³H] DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.1% radioactivity.

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1 50%
NEBuffer 2 75%
NEBuffer 3 75%
NEBuffer 4 **100%**

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 0.13 unit is required to digest 1 µg of substrate DNA in 16 hours.

Notes: BcoDI is an isoschizomer of BsmAI.

Cleavage of mammalian genomic DNA is blocked by some combinations of overlapping CpG methylation.

= Time-Saver™ Qualified (See www.neb.com for details).

CERTIFICATE OF ANALYSIS