

PstI-HF™



1-800-632-7799
info@neb.com
www.neb.com



R3140S 001121014101

R3140S



10,000 units 20,000 U/ml Lot: 0011210

RECOMBINANT Store at -20°C Exp: 10/14

Recognition Site:

5'...CTGCA[▼]G...3'
3'...GACGTC...5'

Note: PstI-HF™ has the same specificity as PstI (NEB #R0140), but it has been engineered for reduced star activity.

Source: An *E. coli* strain that carries the cloned and modified (D91A) PstI gene from *Providencia stuartii* 164 (ATCC 49762)

Supplied in: 250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 0.15% Triton X-100, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:
10X NEBuffer 4.

Reaction Conditions: 1X NEBuffer 4.
Incubate at 37°C.

1X NEBuffer 4:
50 mM potassium acetate
20 mM Tris-acetate
10 mM magnesium acetate
1 mM DTT
pH 7.9 at 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility: Diluent Buffer C
250 mM NaCl, 10 mM Tris-HCl, 0.1 mM EDTA, 1 mM DTT, 0.15% Triton X-100, 200 µg/ml BSA and 50% glycerol (pH 7.4 @ 25°C).

Quality Controls

Ligation: After 100-fold overdigestion with PstI-HF, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 400 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 400 units of enzyme with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.1% radioactivity.

Blue/White Screening Assay: This enzyme has been tested to determine the integrity of the DNA ends produced after digestion with an excess of enzyme. An appropriate vector is digested at a unique site within *lacZ^x* gene with a 10-fold excess of enzyme, ligated, transformed and plated on XGal/IPTG/Amp plates. Successful expression of β-galactosidase is a function of how intact its gene remains after cloning, an intact gene gives rise to

a blue colony, an interrupted gene (i.e. degraded DNA end) gives rise to a white colony. Enzymes must produce fewer than 3% white colonies to be Blue/White Certified.

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1 10%
NEBuffer 2 75%
NEBuffer 3 50%
NEBuffer 4 **100%**

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 0.50 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 80°C for 20 minutes.

Plasmid Cleavage: Number of units required to cleave 1 µg of supercoiled plasmid DNA in one hour: pUC19 = 1 unit, pBR322 = 1 unit, LITMUS = 1 unit.

(see other side)

CERTIFICATE OF ANALYSIS

PstI-HF™



1-800-632-7799
info@neb.com
www.neb.com



R3140S 001121014101

R3140S



10,000 units 20,000 U/ml Lot: 0011210

RECOMBINANT Store at -20°C Exp: 10/14

Recognition Site:

5'...CTGCA[▼]G...3'
3'...GACGTC...5'

Note: PstI-HF™ has the same specificity as PstI (NEB #R0140), but it has been engineered for reduced star activity.

Source: An *E. coli* strain that carries the cloned and modified (D91A) PstI gene from *Providencia stuartii* 164 (ATCC 49762)

Supplied in: 250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 0.15% Triton X-100, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:
10X NEBuffer 4.

Reaction Conditions: 1X NEBuffer 4.
Incubate at 37°C.

1X NEBuffer 4:
50 mM potassium acetate
20 mM Tris-acetate
10 mM magnesium acetate
1 mM DTT
pH 7.9 at 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility: Diluent Buffer C
250 mM NaCl, 10 mM Tris-HCl, 0.1 mM EDTA, 1 mM DTT, 0.15% Triton X-100, 200 µg/ml BSA and 50% glycerol (pH 7.4 @ 25°C).

Quality Controls

Ligation: After 100-fold overdigestion with PstI-HF, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 400 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 400 units of enzyme with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.1% radioactivity.

Blue/White Screening Assay: This enzyme has been tested to determine the integrity of the DNA ends produced after digestion with an excess of enzyme. An appropriate vector is digested at a unique site within *lacZ^x* gene with a 10-fold excess of enzyme, ligated, transformed and plated on XGal/IPTG/Amp plates. Successful expression of β-galactosidase is a function of how intact its gene remains after cloning, an intact gene gives rise to

a blue colony, an interrupted gene (i.e. degraded DNA end) gives rise to a white colony. Enzymes must produce fewer than 3% white colonies to be Blue/White Certified.

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1 10%
NEBuffer 2 75%
NEBuffer 3 50%
NEBuffer 4 **100%**

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 0.50 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 80°C for 20 minutes.

Plasmid Cleavage: Number of units required to cleave 1 µg of supercoiled plasmid DNA in one hour: pUC19 = 1 unit, pBR322 = 1 unit, LITMUS = 1 unit.


(see other side)


CERTIFICATE OF ANALYSIS

Note: Not sensitive to *dam*, *dcm* or mammalian CpG methylation.

New icons (see www.neb.com for details)

 = Time-Saver™ Qualified


 = indicates that the enzyme has been engineered


 = indicates that the enzyme has reduced star activity


Page 2 (R3140)

Note: Not sensitive to *dam*, *dcm* or mammalian CpG methylation.

New icons (see www.neb.com for details)

 = Time-Saver™ Qualified

 = indicates that the enzyme has been engineered

 = indicates that the enzyme has reduced star activity

Page 2 (R3140)