


## Anti-Mouse CD5 FITC

**Catalog Number:** 11-0051

**Also Known As:** Ly-1, Ly-12, Ly-A, Lyl-1

**RUO: For Research Use Only. Not for use in diagnostic procedures.**



Staining of mouse splenocytes with 0.25 µg of Anti-Mouse CD5 FITC (left) or 0.06 µg of Anti-Mouse CD5 PE (right). Appropriate isotype controls were used (open histogram). Total viable cells were used for analysis.

### Product Information

**Contents:** Anti-Mouse CD5 FITC


**REF** **Catalog Number:** 11-0051

**Clone:** 53-7.3


**Concentration:** 0.5 mg/mL


**Host/Isotype:** Rat IgG2a, kappa

**Formulation:** aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 **Temperature Limitation:** Store at 2-8°C. Do not freeze. Light sensitive material.

**LOT** **Batch Code:** Refer to Vial

 **Use By:** Refer to Vial

 **Caution, contains Azide**

### Description

The 53-7.3 monoclonal antibody reacts with mouse CD5, a 67 kDa protein expressed by a majority of thymocytes, mature T cells and a subset of B cells. The expression of CD5 by a small subset of B cells characterizes a developmentally and functionally distinct lineage of B cells called B-1 cells. CD5 is a counter-receptor for CD72 and plays a role in the T-B cell interaction.

### Applications Reported

The 53-7.3 antibody has been reported for use in flow cytometric analysis.

### Applications Tested

The 53-7.3 antibody has been tested by flow cytometric analysis of mouse splenocyte suspensions. This can be used at less than or equal to 0.5 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

### References

Ledbetter JA, Rouse RV, Micklem HS, Herzenberg LA. T cell subsets defined by expression of Lyl-1,2,3 and Thy-1 antigens. Two-parameter immunofluorescence and cytotoxicity analysis with monoclonal antibodies modifies current views. *J Exp Med.* 1980 Aug 1;152 (2):280-95.

Ledbetter JA, Herzenberg LA. Xenogeneic monoclonal antibodies to mouse lymphoid differentiation antigens. *Immunol Rev.* 1979;47:63-90.

### Related Products

11-4321 Rat IgG2a K Isotype Control FITC (eBR2a)

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • [www.eBioscience.com](http://www.eBioscience.com) • [info@eBioscience.com](mailto:info@eBioscience.com)