


## Anti-Human CD40 FITC

**Catalog Number:** 11-0409

**Also known as:** TNFRSF5

**RUO: For Research Use Only. Not for use in diagnostic procedures.**



Staining of normal human peripheral blood cells with Mouse IgG1 K Isotype Control FITC (cat. 11-4714) (blue histogram) or Anti-Human CD40 FITC (purple histogram). Cells in the lymphocyte gate were used for analysis.

### Product Information



**Contents:** Anti-Human CD40 FITC

**Catalog Number:** 11-0409

**Clone:** 5C3

**Concentration:** 5  $\mu$ L (0.5  $\mu$ g)/test

**Host/Isotype:** Mouse IgG1, kappa

**HLDA Workshop:** V B-CD40.4



**Formulation:** aqueous buffer, 0.09% sodium

azide, may contain carrier protein/stabilizer

**Temperature Limitation:** Store at 2-8°C. Do not freeze. Light sensitive material.



**Batch Code:** Refer to vial



**Use By:** Refer to vial

### Description

The 5C3 monoclonal antibody reacts with human CD40, a 45-50 kDa type I transmembrane glycoprotein. CD40 is a member of the TNFR family and is expressed by B lymphocytes, follicular dendritic cells, thymic epithelium, and a subset of peripheral T cells. CD40 regulates B cell development and maturation by inducing Ig isotype-switching and in combination with other signals such as IL-4, protects B cells from surface Ig-induced apoptosis and promotes proliferation. Interaction of CD40 with CD154 (gp39), its ligand on T cells, is important in T-B cell crosstalk and plays a role in costimulation and immune regulation. 5C3 is reported to be used for activation of B cells in *in vitro* functional assays.

### Applications Reported

The 5C3 antibody has been reported for use in flow cytometric analysis.

### Applications Tested

The 5C3 antibody has been pre-titrated and tested by flow cytometric analysis of normal human peripheral blood cells. This can be used at 5  $\mu$ L (0.5  $\mu$ g) per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from  $10^5$  to  $10^8$  cells/test.

### References

Pound JD, Challa A, Holder MJ, Armitage RJ, Dower SK, Fanslow WC, Kikutani H, Paulie S, Gregory CD, Gordon J. Minimal cross-linking and epitope requirements for CD40-dependent suppression of apoptosis contrast with those for promotion of the cell cycle and homotypic adhesions in human B cells. *Int Immunol.* 1999. Jan;11(1):11-20.

Schlossman, S., L. Bloumsell, et al. eds. *Leucocyte Typing V: White Cell Differentiation Antigens.* Oxford University Press. New York. 1995.

### Related Products

11-4714 Mouse IgG1 K Isotype Control FITC (P3.6.2.8.1)

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • [www.ebioscience.com](http://www.ebioscience.com) •  
[info@ebioscience.com](mailto:info@ebioscience.com)

---

## **Anti-Human CD40 FITC**

**Catalog Number:** 11-0409

**Also known as:** TNFRSF5

**RUO: For Research Use Only. Not for use in diagnostic procedures.**

---

12-0209 Anti-Human CD20 PE (2H7)

---

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • [www.ebioscience.com](http://www.ebioscience.com) •  
[info@ebioscience.com](mailto:info@ebioscience.com)