


## Anti-Mouse CD49b (Integrin alpha 2) FITC

Catalog Number: 11-0491

Also Known As: Integrin  $\alpha 2$ , VLA2, ITGA2, DX5

RUO: For Research Use Only



Staining of mouse splenocytes with 0.06  $\mu\text{g}$  of Armenian Hamster IgG Isotype Control FITC (cat. 11-4888) (open histogram) or 0.06  $\mu\text{g}$  of Anti-Mouse CD49b (Integrin  $\alpha 2$ ) FITC (filled histogram). Total viable cells were used for analysis.

### Product Information

Contents: Anti-Mouse CD49b (Integrin alpha 2) FITC


**REF** Catalog Number: 11-0491

Clone: HMa2

Concentration: 0.5 mg/ml


Host/Isotype: Armenian Hamster IgG

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.

**LOT** Batch Code: Refer to Vial

 Use By: Refer to Vial

 Caution, contains Azide

### Description

The HM $\alpha 2$  monoclonal antibody reacts with mouse CD49b, the 150 kDa integrin  $\alpha_2$  subunit. The complex of CD49b non-covalently associated with integrin  $\beta_1$  (CD29), also known as VLA-2, is a receptor for collagen and laminin. This complex is expressed by some CD4<sup>+</sup> T cells, IEL, NK cells, platelets and epithelial cells.

### Applications Reported

The HM $\alpha 2$  antibody has been reported for use in flow cytometric analysis.

### Applications Tested

The HM $\alpha 2$  antibody has been tested by flow cytometric analysis of mouse splenocyte suspensions. This can be used at less than or equal to 0.125  $\mu\text{g}$  per test. A test is defined as the amount ( $\mu\text{g}$ ) of antibody that will stain a cell sample in a final volume of 100  $\mu\text{L}$ . Cell number should be determined empirically but can range from  $10^5$  to  $10^8$  cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

### References

Miyake, S., T. Sakurai, et al. (1994). "Identification of collagen and laminin receptor integrins on murine T lymphocytes." *Eur J Immunol* 24(9): 2000-5.

Noto, K., K. Kato, et al. (1995). "Identification and functional characterization of mouse CD29 with a mAb." *Int Immunol* 7(5): 835-42.

### Related Products

11-4888 Armenian Hamster IgG Isotype Control FITC (eBio299Arm)

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com