


## Anti-Integrin beta 5 FITC

**Catalog Number:** 11-0497

**Also Known As:** Integrin b5, ITGB5

**RUO: For Research Use Only**



Staining of the Chinese Hamster Ovary (CHO) cell line with Mouse IgG1 kappa Isotype Control FITC (cat. 11-4714) (open histogram) or Anti-Integrin beta 5 FITC (filled histogram). Total viable cells were used for analysis.

### Product Information

**Contents:** Anti-Integrin beta 5 FITC


**REF** **Catalog Number:** 11-0497

**Clone:** KN52


**Concentration:** 5 µl (0.125 µg)/test


**Host/Isotype:** Mouse IgG1, kappa

**Formulation:** aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 **Temperature Limitation:** Store at 2-8°C. Do not freeze. Light sensitive material.

**LOT** **Batch Code:** Refer to Vial

 **Use By:** Refer to Vial

 **Caution, contains Azide**

### Description

The KN52 antibody reacts with human integrin beta 5, an approximately 100,000 Mr (reduced) and 95,000 Mr (nonreduced) a member of the integrin family. Only a single alpha chain, the alpha v subunit, associates with the integrin beta 5 to form the vitronectin receptor complex. Integrin alpha v/beta 5 complex also binds to the basic domain of Tat (he sequence RKKRRQRRR). The integrin beta 5 is found on carcinoma cell lines, hepatoma and fibroblast cell lines, and is absent from lymphoblastoid cells and platelets.

The KN52 monoclonal antibody has been shown to inhibit migration.

### Applications Reported

This KN52 antibody has been reported for use in flow cytometric analysis.

### Applications Tested

This KN52 antibody has been pre-titrated and tested by flow cytometric analysis of Chinese Hamster Ovary (CHO), A549 and normal human peripheral blood cells. This can be used at 5 µl (0.125 µg)/per test. A test is defined as the amount (µg)/test of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test.

### References

Sangaletti S, Di Carlo E, Gariboldi S, Miotti S, Cappetti B, Parenza M, Rumio C, Brekken RA, Chiodoni C, Colombo MP. Macrophage-derived SPARC bridges tumor cell-extracellular matrix interactions toward metastasis. *Cancer Res.* 2008 Nov 1;68(21):9050-9. (**KN52**, FA, PubMed)

Lai CF, Feng X, Nishimura R, Teitelbaum SL, Avioli LV, Ross FP, Cheng SL. Transforming growth factor-beta up-regulates the beta 5 integrin subunit expression via Sp1 and Smad signaling. *J Biol Chem.* 2000 Nov 17;275(46):36400-6.

Sugiyama M, Speight PM, Prime SS, Watt FM. Comparison of integrin expression and terminal differentiation capacity in cell lines derived from oral squamous cell carcinomas. *Carcinogenesis.* 1993 Oct;14(10):2171-6.

### Related Products

11-4714 Mouse IgG1 K Isotype Control FITC

