


Anti-Human CD56 (NCAM) FITC

Catalog Number: 11-0569

Also Known As:NCAM, Neural cell adhesion molecule

RUO: For Research Use Only



Staining of normal human peripheral blood cells with Anti-Human CD94 PE (cat. 12-0949) and Anti-Human CD56 (NCAM) FITC. Cells in the lymphocyte gate were used for analysis.

Product Information

Contents: Anti-Human CD56 (NCAM) FITC

REF Catalog Number: 11-0569
Clone: MEM188 (MEM-188)
Concentration: 5 μl (1 μg)/test
Host/Isotype: Mouse IgG2a, κ
HLDA Workshop: VI NK26

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light

sensitive material.

Batch Code: Refer to Vial

Use By: Refer to Vial

Caution, contains Azide

Description

The MEM188 monoclonal antibody reacts with human CD56, also known as Neural Cell Adhesion Molecule (NCAM). CD56 is a highly glycosylated transmembrane molecule expressed by neurons and plays a role in the homotypic adhesion of neural cells. In the hematopoietic system, CD56 is expressed on NK cells and a subset of T cells referred to as NKT cells.

Applications Reported

The MEM188 (MEM-188) antibody has been reported for use in flow cytometric analysis.

Applications Tested

This MEM188 (MEM-188) antibody has been pre-titrated and tested by flow cytometric analysis of human peripheral blood leukocytes. This can be used at 5 μ l (1 μ g)/per test. A test is defined as the amount (μ g)/test of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test.

References

Kishimoto, T., A.E.G., von dem Borne, et al. eds. 1998. Leucocyte Typing VI: White Cell Differentiation Antigens. Garland Publishing, Inc. London.

Related Products

11-4724 Mouse IgG2a K Isotype Control FITC