


Anti-Mouse CD69 FITC

Catalog Number: 11-0691

Also known as: Very Early Activation Antigen, VEA

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of unstimulated (left) or overnight ConA-stimulated (right) C57BL/6 mouse splenocytes with 0.25 ug of Armenian Hamster IgG Isotype Control FITC (cat. 11-4888) (blue histogram) or 0.25 ug of Anti-Mouse CD69 FITC (purple histogram). Total viable cells were analyzed.

Product Information



Contents: Anti-Mouse CD69 FITC

Catalog Number: 11-0691

Clone: H1.2F3

Concentration: 0.5 mg/mL

Host/Isotype: Armenian Hamster IgG



Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.

Batch Code: Refer to vial

Use By: Refer to vial

Caution, contains Azide



Description

The H1.2F3 monoclonal antibody reacts with mouse CD69, also known as very early activation antigen (VEA). CD69 is approximately 35 kDa and is expressed on the surface as a disulfide-linked dimer. While a small subset of lymphocytes in the thymus, spleen and lymph nodes express this antigen, activation of both T and B cells rapidly upregulates the surface expression of CD69, suggesting a role for CD69 in lymphocyte development and activation.

Applications Reported

The H1.2F3 antibody has been reported for use in flow cytometric analysis.

Applications Tested

The H1.2F3 antibody has been tested by flow cytometric analysis of resting and activated mouse splenocyte suspensions. This can be used at less than or equal to 0.5 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Yokoyama, W. M., F. Koning, et al. (1988). "Characterization of a cell surface-expressed disulfide-linked dimer involved in murine T cell activation." *J Immunol* 141(2): 369-76.

Related Products

11-4888 Armenian Hamster IgG Isotype Control FITC (eBio299Arm)

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com