

Anti-Mouse/Rat TNF alpha FITC

Catalog Number: 11-7423

Also known as: Tumor Necrosis Factor alpha RUO: For Research Use Only. Not for use in diagnostic procedures.



Mouse splenocytes were unstimulated (left) or stimulated for 5 hours with Cell Stimulation Cocktail (plus protein transport inhibitors) (cat. 00-4975) (right), then intracellularly stained with Anti-Mouse CD4 eFluor® 450 (cat. 48-0041) and 0.125 ug of Anti-Mouse/Rat TNF alpha FITC. Total viable cells, as determined by Fixable Viability Dye eFluor® 780, were used for analysis.

Product Information

Contents: Anti-Mouse/Rat TNF alpha FITC
Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

REF
Catalog Number: 11-7423

Clone: TN3-19
Concentration: ug size: 0.5 mg/mL; test size: 5 uL (0.125 ug)/test

Host/Isotype: Armenian Hamster IgG
Batch Code: Refer to vial

Image: Solution of the section of th

Description

The TN3-19.12 monoclonal antibody reacts with mouse, rat, and rabbit tumor necrosis factor-alpha (TNF alpha), but not with human TNF alpha. The TN3-19.12 antibody is a neutralizing antibody. Mouse TNF alpha is a 17 kDa factor produced by macrophages, monocytes, neutrophils, CD4+ T cells and NK cells. A 26 kDa form of TNF alpha is expressed as a membrane-bound molecule. TNF alpha is cytolytic and plays an important role in immune regulation. Dimers and trimers of TNF alpha have been observed.

Applications Reported

The TN3-19.12 antibody has been reported for use in intracellular staining for flow cytometric analysis, cytokine neutralization, and ELISA. Fluorochrome conjugated TN3-19.12 is recommended for use in intracellular staining and flow cytometry. Functional Grade purified TN3-19.12, cat. 16-7423, is recommended for use in functional assays. For ELISPOT capture, the alternative clone, 1F3F3D4 (cat # 16-7325), is recommended.

Applications Tested

This TN3-19.12 antibody has been tested by intracellular staining and flow cytometric analysis of stimulated mouse splenocytes. It is offered in 2 formats:

- µg size: can be used at less than or equal to 0.25 µg per test.
- test size: can be used at 5 µL (0.125 µg) per test.

A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test.

References

Finkelman, F.D., and S.C. Morris. Development of an assay to measure in vivo cytokine production in the mouse. Int. Immunology. 1999. 11: 1811-1818.

Finkelman, F., S. Morris, T. Orekhova, and D. Sehy. 2003. The In Vivo Cytokine Capture Assay for measurement of

Not for further distribution without written consent. Copyright © 2000-2012 eBioscience, Inc. Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com • info@ebioscience.com



Anti-Mouse/Rat TNF alpha FITC

Catalog Number: 11-7423 Also known as: Tumor Necrosis Factor alpha RUO: For Research Use Only. Not for use in diagnostic procedures.

cytokine production in the mouse. In Current Protocols in Immunology. Unit 6.28. J. Coligan, A. Kruisbeek, D. Margulies, E. Shevach, and W. Strober, eds. John Wiley and Sons, New York.

Takahashi S, Kapas L, Fang J, Krueger JM. An anti-tumor necrosis factor antibody suppresses sleep in rats and rabbits. Brain Res. 1995. 690: 241-44.

Rabinovici R, Bugelski PJ, Esser KM, Hillegass LM, Griswold DE, Vernick J, Feuerstein G.Tumor necrosis factoralpha mediates endotoxin-induced lung injury in platelet activating factor-primed rats. Macol Exp Ther. 1993. 267: 1550-57.

Merrick BA, He CY, Craig WA, Clark GC, Corsini E, Rosenthal GJ, Mansfield BK, Selkirk JK. Two dimensional gel electrophoresis of cellular and secreted proteins from rat alveolar macrophages after lipopolysaccharide treatment. Appl Theor Electrophor. 1992. 2: 177-87.

Sheehan KC, Ruddle NH, Schreiber RD. Generation and characterization of hamster monoclonal antibodies that neutralize murine tumor necrosis factors. J Immunol. 1989. 142: 3884-93.

Related Products

00-4975 Cell Stimulation Cocktail (plus protein transport inhibitors) (500X) 11-4888 Armenian Hamster IgG Isotype Control FITC (eBio299Arm) 14-8021 Mouse IL-2 Recombinant Protein 14-8041 Mouse IL-4 Recombinant Protein 16-0032 Anti-Mouse CD3 Functional Grade Purified (17A2) 16-0281 Anti-Mouse CD28 Functional Grade Purified (37.51) 48-0041 Anti-Mouse CD4 eFluor® 450 (GK1.5) 65-0865 Fixable Viability Dye eFluor® 780 88-8823 Fixation & Permeabilization Buffers