


Anti-Mouse/Rat CD29 (Integrin beta 1) Purified

Catalog Number: 14-0291

Also Known As: Fibronectin Receptor Beta

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of C57Bl/6 thymocytes with 0.5 ug of Armenian Hamster IgG Isotype Control Purified (cat. 14-4888) (open histogram) or 0.5 ug of Anti-Mouse/Rat CD29 Purified (filled histogram) followed by Anti-Armenian Hamster IgG Biotin (cat. 13-4113) and Streptavidin PE (cat. 12-4317). Total viable cells were used for analysis.

Product Information

Contents: Anti-Mouse/Rat CD29 (Integrin beta 1) Purified

REF **Catalog Number:** 14-0291

Clone: eBioHMb1-1 (HMb1-1)

Concentration: 0.5 mg/mL

Host/Isotype: Armenian Hamster IgG

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer



Temperature Limitation: Store at 2-8°C.



Batch Code: Refer to Vial



Use By: Refer to Vial



Contains sodium azide

Description

The eBioHMb1-1 monoclonal antibody reacts with mouse and rat CD29 (integrin beta 1), a 110-120 kDa member of the beta integrin family expressed by leukocytes, endothelial, smooth muscle and epithelial cells. CD29 binds non-covalently with the alpha integrins CD49a-f to form the VLA-1 through VLA-6 complexes, as well as with CD51. These alpha-beta integrin heterodimers are capable of mediating a variety of cellular responses including adhesion, trafficking, proliferation and differentiation. All integrins which include CD29 bind to extracellular matrix proteins including collagen, laminin, fibronectin and vitronectin, whereas some CD29-containing integrins can also interact with cellular receptors such as VCAM-1 and MadCAM-1.

Applications Reported

This eBioHMb1-1 (HMb1-1) antibody has been reported for use in flow cytometric analysis, and immunoprecipitation. (Please use Functional Grade purified eBioHMb1-1 (HMb1-1), cat. 16-0291, in functional assays.)

Applications Tested

This eBioHMb1-1 (HMb1-1) antibody has been tested by flow cytometric analysis of mouse spleen, thymus and bone marrow cell suspensions. This can be used at less than or equal to 1 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Noto K, Kato K, Okumura K, Yagita H. Identification and functional characterization of mouse CD29 with a mAb. *Int Immunol.* 1995 May;7(5):835-42. (**HMBeta1-1**, mAb development, IP, FA, PubMed)

Ridger VC, Wagner BE, Wallace WA, Hellewell PG. Differential effects of CD18, CD29, and CD49 integrin subunit inhibition on neutrophil migration in pulmonary inflammation. *J Immunol.* 2001 Mar 1;166(5):3484-90. (**HMBeta1-1**, FA, PubMed)

Related Products

11-4111 Anti-Armenian Hamster IgG FITC

11-4317 Streptavidin FITC

12-4317 Streptavidin PE

13-4113 Anti-Armenian Hamster IgG Biotin (Polyclonal)

14-4888 Armenian Hamster IgG Isotype Control Purified (eBio299Arm)

17-4317 Streptavidin APC

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com