


Anti-Mouse Siglec H Purified

Catalog Number: 14-0333

Also known as: Sialic Acid-Binding Immunoglobulin-Like Lectin H

RUO: For Research Use Only. Not for use in diagnostic procedures.

Product Information

 **Contents:** Anti-Mouse Siglec H Purified
Catalog Number: 14-0333
Clone: eBio440c
Concentration: 0.5 mg/mL
Host/Isotype: Rat IgG2b



Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer
Temperature Limitation: Store at 2-8°C.
Batch Code: Refer to vial
Use By: Refer to vial
Contains sodium azide

Description

The monoclonal antibody eBio440c recognizes Siglec-H, a protein exclusively found on pDC (plasmacytoid dendritic cells) or type I IFN-producing cells (IPC) in the naïve mouse. Mouse IPC are typically PDCA+, CD11c+, CD11b-, B220+, and Ly-6C+, and are quick to respond to viruses. Siglec-H is a transmembrane protein of the Ig superfamily that like CD33 have been shown to bind sialic acid but lacks the characteristic cytoplasmic ITIM domain (immunoreceptor tyrosine based inhibitory motif). To overcome the lack of a cytoplasmic domain, Siglec-H associates with DAP12 thereby allowing for signal transduction.

The eBio440c antibody has been shown to inhibit pDC function (inhibits IFN α secretion in response to CpG).

It has been observed that some mouse strains (such as SJL) have higher percentages of pDCs compared to C57BL/6.

Applications Tested

This eBio440c antibody has been tested by blocking staining of the direct conjugate of eBio440c. This can be used at less than or equal to 2 μ g per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test.

It is recommended that direct conjugates such as cat. 11-0333 be used rather than purified eBio440c antibody when performing flow cytometric analysis.

References

Kreisel FH, Blasius A, Kreisel D, Colonna M, Cella M. Interferon-producing cells develop from murine CD31(high)/Ly6C(-) marrow progenitors. *Cell Immunol.* 2006 Aug;242(2):91-8.

Blasius AL, Cella M, Maldonado J, Takai T, Colonna M. Siglec-H is an IPC-specific receptor that modulates type I IFN secretion through DAP12. *Blood.* 2006 Mar 15;107(6):2474-6.(440c, FC, PubMed)

Blasius AL, Giurisato E, Cella M, Schreiber RD, Shaw AS, Colonna M. Bone marrow stromal cell antigen 2 is a specific marker of type I IFN-producing cells in the naive mouse, but a promiscuous cell surface antigen following IFN stimulation. *J Immunol.* 2006 Sep 1;177(5):3260-5.(440c, FC, PubMed)

Blasius A, Vermi W, Krug A, Facchetti F, Cella M, Colonna M. A cell-surface molecule selectively expressed on murine natural interferon alpha producing cells that blocks secretion of interferon alpha. *Blood.* 2004;103:4201-4206 (440c, FC, IH/F, FA PubMed)

Related Products

11-4811 Anti-Rat IgG FITC
12-3172 Anti-Mouse CD317 (BST2, PDCA-1) PE (eBio927)
14-4031 Rat IgG2b K Isotype Control Purified

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com

Anti-Mouse Siglec H Purified

Catalog Number: 14-0333

Also known as: Sialic Acid-Binding Immunoglobulin-Like Lectin H

RUO: For Research Use Only. Not for use in diagnostic procedures.

Legal

Pat. No. pending

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com