


Anti-Human CD48 Purified

Catalog Number: 14-0489

Also Known As: BCM1, BCM 1, B-cell membrane protein, SLAMF2, SLAM F2, BLAST1

RUO: For Research Use Only



Staining of normal human peripheral blood mononuclear cells with 0.5 µg of Mouse IgG1 κ Isotype Control Purified (cat. 14-4714) (open histogram) or 0.5 µg of Anti-Human CD48 Purified (filled histogram) followed by F(ab')₂ Anti-Mouse IgG PE (cat. 12-4012). Cells in the lymphocyte gate were used for analysis.

Product Information

Contents: Anti-Human CD48 Purified

REF Catalog Number: 14-0489

Clone: eBio156-4H9 (156-4H9)

Concentration: 0.5 mg/ml

Host/Isotype: Mouse IgG1

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer



Temperature Limitation: Store at 2-8°C.



Batch Code: Refer to Vial



Use By: Refer to Vial



Caution, contains Azide

Description

The eBio156-4H9 monoclonal antibody reacts with the human CD48 antigen, also known as B cell activation marker 1, SLAM family member 2, Blast-1 (human), and OX-45 (rat). CD48, a member of the SLAM family and Ig superfamily, is a 45 kDa GPI-linked glycoprotein expressed on the majority of hematopoietic cells. CD2 is the low affinity ligand for CD48. Recent publications have reported differential expression in mice of members of the SLAM family including CD48, CD150, and CD244 among functionally distinct bone marrow hematopoietic progenitors providing a useful tool for prediction of the primitiveness of hematopoietic progenitors. Expression of CD48 in hematopoietic cells is found on many leukocytes with little to none on granulocytes and absent on platelets and erythrocytes. Expression increases on B cells with activation.

Applications Reported

This eBio156-4H9 (156-4H9) antibody has been reported for use in flow cytometric analysis.

Applications Tested

This eBio156-4H9 (156-4H9) antibody has been tested by flow cytometric analysis of human peripheral blood cells. This can be used at less than or equal to 1 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Yokoyama S, Staunton D, Fisher R, Amiot M, Fortin JJ, Thorley-Lawson DA. Expression of the Blast-1 activation/adhesion molecule and its identification as CD48. *J Immunol.* 1991 Apr 1;146(7):2192-200.

Gonzalez-Alvaro I, Dominguez-Jimenez C, Ortiz AM, Nunez-Gonzalez V, Roda-Navarro P, Fernandez-Ruiz E, Sancho D, Sanchez-Madrid F. Interleukin-15 and interferon-gamma participate in the cross-talk between natural killer and monocytic cells required for tumour necrosis factor production. *Arthritis Res Ther.* 2006;8(4):R88. (156-4H9, FC, PubMed)

Stonehouse TJ, Woodhead VE, Herridge PS, Ashrafian H, George M, Chain BM, Katz DR. Molecular characterization of U937-dependent T-cell co-stimulation. *Immunology.* 1999 Jan;96(1):35-47. (156-4H9, FC, PubMed)

Related Products

11-4011 Anti-Mouse IgG FITC

14-0481 Anti-Mouse CD48 Purified (HM48-1)

14-4714 Mouse IgG1 K Isotype Control Purified

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com