


Anti-Integrin beta 5 Purified

Catalog Number: 14-0497

Also Known As: Integrin b5, ITGB5

RUO: For Research Use Only



Staining of Chinese Hamster Ovary (CHO) cell line with 0.25 ug of Mouse IgG1kappa Isotype Control Purified (cat. 14-4714) (open histogram) or 0.25 ug of Anti-Integrin beta 5 Purified (filled histogram) followed by Anti-Mouse IgG FITC (cat. 11-4011). Total viable cells were used for analysis.

Product Information

Contents: Anti-Integrin beta 5 Purified

REF **Catalog Number:** 14-0497

Clone: KN52

Concentration: 0.5 mg/ml

Host/Isotype: Mouse IgG1, kappa

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer



Temperature Limitation: Store at 2-8°C.



Batch Code: Refer to Vial



Use By: Refer to Vial



Caution, contains Azide

Description

The KN52 antibody reacts with human integrin beta 5, an approximately 100,000 Mr (reduced) and 95,000 Mr (nonreduced) a member of the integrin family. Only a single alpha chain, the alpha v subunit, associates with the integrin beta 5 to form the vitronectin receptor complex. Integrin alpha v/beta 5 complex also binds to the basic domain of Tat (he sequence RKKRRQRRR). The integrin beta 5 is found on carcinoma cell lines, hepatoma and fibroblast cell lines, and is absent from lymphoblastoid cells and platelets.

The KN52 monoclonal antibody has been shown to inhibit migration.

Applications Reported

This KN52 antibody has been reported for use in flow cytometric analysis. (Please use Functional Grade purified KN52, cat. 16-0497, in functional assays.)

Applications Tested

This KN52 antibody has been tested by flow cytometric analysis of Chinese Hamster Ovary (CHO), A549 and normal human peripheral blood cells. This can be used at less than or equal to 0.25 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Sangaletti S, Di Carlo E, Gariboldi S, Miotti S, Cappetti B, Parenza M, Rumio C, Brekken RA, Chiodoni C, Colombo MP. Macrophage-derived SPARC bridges tumor cell-extracellular matrix interactions toward metastasis. *Cancer Res.* 2008 Nov 1;68(21):9050-9. (**KN52**, FA, PubMed)

Lai CF, Feng X, Nishimura R, Teitelbaum SL, Avioli LV, Ross FP, Cheng SL. Transforming growth factor-beta up-regulates the beta 5 integrin subunit expression via Sp1 and Smad signaling. *J Biol Chem.* 2000 Nov 17;275(46):36400-6.

Sugiyama M, Speight PM, Prime SS, Watt FM. Comparison of integrin expression and terminal differentiation capacity in cell lines derived from oral squamous cell carcinomas. *Carcinogenesis.* 1993 Oct;14(10):2171-6.

Related Products

11-4011 Anti-Mouse IgG FITC

11-4317 Streptavidin FITC

12-4317 Streptavidin PE

13-4013 Anti-Mouse IgG Biotin (Polyclonal)

14-4714 Mouse IgG1 K Isotype Control Purified
17-4317 Streptavidin APC

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com