


Anti-Mouse MHC Class II (I-A) Purified

Catalog Number: 14-5322

Also Known As: MHC II, IA

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of C57BL/6 splenocytes with 0.25 ug of Rat IgG2b K Isotype Control Purified (cat. 14-4031) (open histogram) or 0.25 ug of Anti-Mouse MHC Class II (I-A) Purified (filled histogram) followed by Anti-Rat IgG FITC (cat. 11-4811). Cells in the lymphocyte gate were used for analysis.

Product Information

Contents: Anti-Mouse MHC Class II (I-A) Purified

REF **Catalog Number:** 14-5322

Clone: NIMR-4

Concentration: 0.5 mg/mL

Host/Isotype: Rat IgG2b

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer



Temperature Limitation: Store at 2-8°C.



Batch Code: Refer to Vial



Use By: Refer to Vial



Caution, contains Azide

Description

The monoclonal antibody NIMR-4 reacts specifically with a non-polymorphic I-A-encoded epitope on MHC Class II antigens. Class II antigens are predominantly expressed on antigen-presenting cells including B lymphocytes, macrophages, dendritic cells and certain epithelial cells. MHC Class II molecules are important for the selection of the TCR repertoire during the processes of positive and negative selection. MHC Class II expression is required for the positive selection of CD4+ T cells.

Applications Reported

This NIMR-4 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This NIMR-4 antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to 1 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Mielenz, D., C. Vetterman, M. Hampel, C. Lang, A. Avramidou, M. Karas, and H. Jack. 2005. Lipid rafts associate with intracellular B cell receptors and exhibit a B cell stage-specific protein composition. *J. Immunol.* 174: 3508-3517. (NIMR-4, FC, FA, PubMed)

Related Products

11-4811 Anti-Rat IgG FITC

17-4317 Streptavidin APC

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com