


Anti-Mouse/Rat Foxp3 Purified

Catalog Number: 14-5773

Also Known As: Forkhead Box P3, Scurfin, JM2, Treg

RUO: For Research Use Only



Immunoblotting of BALB/c spleen (Lane A), BALB/c thymus (Lane B), or BALB/c bone marrow (Lane C) lysate with 2 µg/mL of Anti-Mouse/Rat Foxp3 Purified and revealed with Anti-Rat IgG HRP . Foxp3 is detected at ~50 kDa.

Product Information

Contents: Anti-Mouse/Rat Foxp3 Purified

REF Catalog Number: 14-5773

Clone: FJK-16s

Concentration: 0.5 mg/ml

Host/Isotype: Rat IgG2a, κ

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 Temperature Limitation: Store at 2-8°C.

 Batch Code: Refer to Vial

 Use By: Refer to Vial

 Caution, contains Azide

Description

The FJK-16s antibody reacts with mouse, rat, dog, porcine, bovine and cat Foxp3 also known as FORKHEAD BOX P3, SCURFIN, and JM2; cross reactivity of this antibody to other proteins has not been determined. Foxp3, a 49-55 kDa protein, is a member of the forkhead/winged-helix family of transcriptional regulators, and was identified as the gene defective in 'scurfy' (sf) mice. Constitutive high expression of foxP3 mRNA has been shown in CD4+CD25+ regulatory T cells (Treg cells), and ectopic expression of foxp3 in CD4+CD25- cells imparts a Treg phenotype in these cells.

Immunoblotting with FJK-16s antibody has mapped the epitope to amino acids 75-125 of the mouse Foxp3 protein. In the human, this region has been shown to be alternatively spliced at the mRNA level. Both the alternatively-spliced and non-spliced isoforms are present in the CD4+CD25+ subset of lymphocytes. Preliminary RT-PCR experiments have not revealed this alternatively-spliced isoform in mouse splenocytes, suggesting different gene regulation in the mouse and human.

Applications Reported

This FJK-16s antibody has been reported for use in immunoblotting (WB) and immunohistological staining of frozen and paraffin embedded tissue sections. For additional information, please see <http://www.ebioscience.com/ebioscience/Foxp3FAQs.htm>.

Applications Tested

This FJK-16s antibody has been tested by immunoblotting (WB) (1-5µg/ml) of mouse splenocytes.

References

Gerner W, Stadler M, Hammer SE, Klein D, Saalmüller A. Sensitive detection of Foxp3 expression in bovine lymphocytes by flow cytometry. *Vet Immunol Immunopathol.* 2010 Nov 15;138(1-2):154-8 (FJK-16s, porcine FC, PubMed)

Käser T, Gerner W, Hammer SE, Patzl M, Saalmüller A. Detection of Foxp3 protein expression in porcine T lymphocytes. *Vet Immunol Immunopathol.* 2008 Sep 15;125(1-2):92-101 (FJK-16s, IC flow porcine, PubMed)

Habicht A, Dada S, Jurewicz M, Fife BT, Yagita H, Azuma M, Sayegh MH, Guleria I. A link between PDL1 and T regulatory cells in fetomaternal tolerance. *J Immunol.* 2007 Oct 15;179(8):5211-9. (FJK-16s, IH/F, PubMed)

Heimesaat MM, Fischer A, Siegmund B, Kupz A, Niebergall J, Fuchs D, Jahn HK, Freudenberg M, Loddenkemper C, Batra A, Lehr HA, Liesenfeld O, Blaut M, Göbel UB, Schumann RR, Bereswill S. Shift towards pro-inflammatory bacteria aggravates acute murine colitis via Toll-like receptors 2 and 4. *PLoS ONE.* 2007 Jul 25;2(7):e662. (FJK-16s, IHC paraffin, PubMed)

Biller BJ, Elmslie RE, Burnett RC, Avery AC, Dow SW. Use of FoxP3 expression to identify regulatory T cells in healthy dogs and dogs with cancer. *Vet Immunol Immunopathol.* 2007 Mar 15;116(1-2):69-78. (FJK-16s, IC Flow in canine, PubMed)

Leithäuser F, Meinhardt-Krajina T, Fink K, Wotschke B, Möller P, Reimann J. Foxp3-expressing CD103⁺ regulatory T cells accumulate in dendritic cell aggregates of the colonic mucosa in murine transfer colitis. *Am J Pathol.* 2006 Jun;168(6):1898-909. (FJK-16s, IHC paraffin, PubMed)

Kohm AP, McMahon JS, Podojil JR, Begolka WS, Degutes M, Kasprowicz DJ, Ziegler SF, Miller SD. Cutting Edge: Anti-CD25 Monoclonal Antibody Injection Results in the Functional Inactivation, Not Depletion, of CD4+CD25⁺ T Regulatory Cells. *J Immunol.* 2006 Mar 15;176(6):3301-5. (FJK-16s; intracellular staining and IH/F, PubMed)

Suvas S, Azkur AK, Rouse BT. Qa-1b and CD94-NKG2a interaction regulate cytolytic activity of herpes simplex virus-specific memory CD8⁺ T cells in the latently infected trigeminal ganglia. *J Immunol.* 2006 Feb 1;176(3):1703-11. (FJK-16s, Immunofluorescence, PubMed)

Fields ML, Hondowicz BD, Metzgar MH, Nish SA, Wharton GN, Picca CC, Caton AJ, Erikson J. CD4+CD25⁺ Regulatory T cells inhibit the maturation but not the initiation of an autoantibody response. *J Immunol.* 2005 Oct 1;175(7):4255-64. (FJK-16s, IC Flow, PubMed)

Ko K, Yamazaki S, Nakamura K, Nishioka T, Hirota K, Yamaguchi T, Shimizu J, Nomura T, Chiba T, Sakaguchi S. Treatment of advanced tumors with agonistic anti-GITR mAb and its effects on tumor-infiltrating Foxp3⁺CD25⁺CD4⁺ regulatory T cells. *J Exp Med.* 2005 Oct 3;202(7):885-91. (FJK-16s, IC Flow, PubMed)

Aswad F, Kawamura H, Dennert G. High Sensitivity of CD4+CD25⁺ Regulatory T Cells to Extracellular Metabolites Nicotinamide Adenine Dinucleotide and ATP: A Role for P2X7 Receptors. *J Immunol.* 2005 Sep 1;175(5):3075-83. (FJK-16s, IC Flow, PubMed)

Beyersdorf N, Gaupp S, Balbach K, Schmidt J, Toyka KV, Lin CH, Hanke T, Hünig T, Kerckau T, Gold R. Selective targeting of regulatory T cells with CD28 superagonists allows effective therapy of experimental autoimmune encephalomyelitis. *J Exp Med.* 2005 Aug 1;202(3):445-55. (FJK-16s, IC Flow in Rat, PubMed)

Fontenot JD, Rasmussen JP, Williams LM, Dooley JL, Farr AG, Rudensky AY. Regulatory T cell lineage specification by the forkhead transcription factor foxp3. *Immunity.* 2005 Mar;22(3):329-41.

Hori S, Nomura T, Sakaguchi S. Control of regulatory T cell development by the transcription factor Foxp3. *Science.* 2003 Feb 14;299 (5609):1057-61.

Related Products

00-5523 Foxp3 Staining Buffer Set

14-4321 Rat IgG2a K Isotype Control Purified

72-5776 Anti-Human Foxp3 Staining Set PE (PCH101 Set)

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com