


## Anti-Human CD314 (NKG2D) Purified

Catalog Number: 14-5878

Also Known As: KLRK1

RUO: For Research Use Only



Staining of normal human peripheral blood cells with 0.06 µg of Mouse IgG1 κ Isotype Control Purified (cat. 14-4714) (open histogram) or 0.06 µg of Anti-Human CD314 (NKG2D) Purified (filled histogram) followed by Anti-Mouse IgG FITC (cat. 11-4011). Cells in the lymphocyte gate were used for analysis.

### Product Information

Contents: Anti-Human CD314 (NKG2D) Purified

**REF** Catalog Number: 14-5878

Clone: 1D11

Concentration: 0.5 mg/ml

Host/Isotype: Mouse IgG1

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer



Temperature Limitation: Store at 2-8°C.



Batch Code: Refer to Vial



Use By: Refer to Vial



Caution, contains Azide

### Description

The 1D11 monoclonal antibody reacts with the human NKG2D, a 42 kDa lectin-like molecule expressed by NK cells, γδ T cells, some CD4<sup>+</sup> and CD8<sup>+</sup> T cells. Human NKG2D forms complexes with DAP10, a membrane adaptor protein, and has the ability to costimulate multiple NK activation receptors. The counter-receptor for human NKG2D has been identified as MICA/MICB expressed on epithelial tumors from lung, breast, kidney, ovary, prostate and colon carcinoma. 5C6 and 1D11 block binding of soluble MICA to γδ TCR T cell clones and inhibit lysis by these cells. 5C6 and 1D11 induced NKG2D function of redirected lysis of FcReceptor bearing P815 cells.

### Applications Reported

The 1D11 antibody has been reported for use in flow cytometric analysis, immunoprecipitation, and immunohistochemical staining. It has also been reported in inhibition of ligand binding in *in vitro* assays. Please use Functional Grade purified 1D11 in functional assays.)

### Applications Tested

The 1D11 antibody has been tested by flow cytometric analysis of human peripheral blood leukocytes. This can be used at less than or equal to 0.125 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

### References

Veronika Groh, Alexander Steinle, Stefan Bauer, and Thomas Spies. 1998. Recognition of Stress-Induced MHC Molecules by Intestinal Epithelial T Cells. *Science*. 279:1737-1740.

Stefan Bauer, Veronika Groh, Jun Wu, Alexander Steinle, Joseph H. Phillips, Lewis L. Lanier, and Thomas Spies. 1999. Activation of NK Cells and T Cells by NKG2D, a Receptor for Stress-Inducible MICA. *Science*. 285: 727-729.

Groh V, Bruhl A, et al. 2003. Stimulation of T cell autoreactivity by anomalous expression of NKG2D and its MIC ligands in rheumatoid arthritis. *Proc Natl Acad Sci U S A*. 100(16):9452-7, (IHC frozen, PubMed)

Roberts AI, Lee L, et al. 2001. NKG2D receptors induced by IL-15 costimulate CD28-negative effector CTL in the tissue microenvironment. *J Immunol*. 167(10):5527-30. (activation, PubMed)

### Related Products

11-4011 Anti-Mouse IgG FITC

11-4317 Streptavidin FITC

12-4317 Streptavidin PE

13-4013 Anti-Mouse IgG Biotin (Polyclonal)

14-4714 Mouse IgG1 K Isotype Control Purified

14-5879 Anti-Human CD314 (NKG2D) Purified (5C6)

17-4317 Streptavidin APC

---

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • [www.eBioscience.com](http://www.eBioscience.com) • [info@eBioscience.com](mailto:info@eBioscience.com)