


Anti-Mouse CD252 (OX40 Ligand) Purified

Catalog Number: 14-5905

Also Known As: CD134L, CD134 Ligand, TNFSF4, gp34

RUO: For Research Use Only. Not for use in diagnostic procedures.



Surface staining of Anti-Mouse IgM and Anti-Mouse CD40-stimulated splenocytes with Anti-Mouse CD252 (OX40 Ligand) PE. Appropriate isotype controls were used (open histogram). Total viable cells were used for analysis.

Product Information

Contents: Anti-Mouse CD252 (OX40 Ligand) Purified

REF **Catalog Number:** 14-5905

Clone: RM134L

Concentration: 0.5 mg/mL

Host/Isotype: Rat IgG2b, kappa

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer



Temperature Limitation: Store at 2-8°C.



Batch Code: Refer to Vial



Use By: Refer to Vial



Caution, contains Azide

Description

The RM134L monoclonal antibody reacts with mouse CD252 also known as OX-40 Ligand, a member of the TNF superfamily. OX-40L is induced on mouse splenic B cells stimulated with a combination of anti-IgM and anti-CD40. Neither resting nor activated mouse T cells express OX-40L. The interaction of OX-40 (CD134) with OX-40L has been implicated in T cell-dependent humoral response. RM134L inhibits the costimulatory activity of mouse OX-40L transfected cells for anti-CD3-stimulated T cell proliferation.

Applications Reported

This RM134L antibody has been reported for use in flow cytometric analysis, immunoprecipitation, and immunohistology staining of frozen tissue sections. (Please use Functional Grade purified RM134L, cat. 16-5905, in functional assays.)

Applications Tested

The RM134L antibody has been tested by flow cytometric analysis of activated mouse splenocyte suspensions and can be used at less than or equal to 0.5 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Akiba H, Oshima H, Takeda K, Atsuta M, Nakano H, Nakajima A, Nohara C, Yagita H, Okumura K. CD28-independent costimulation of T cells by OX40 ligand and CD70 on activated B cells. *J Immunol.* 1999 Jun 15;162(12):7058-66. (RM134L, FA blocking)

Akiba H, Miyahira Y, Atsuta M, Takeda K, Nohara C, Futagawa T, Matsuda H, Aoki T, Yagita H, Okumura K. Critical contribution of OX40 ligand to T helper cell type 2 differentiation in experimental leishmaniasis. *Exp Med.* 2000 Jan 17;191(2):375-80.

Tsukada N, Akiba H, Kobata T, Aizawa Y, Yagita H, Okumura K. Blockade of CD134 (OX40)-CD134L interaction ameliorates lethal acute graft-versus-host disease in a murine model of allogeneic bone marrow transplantation. *Blood.* 2000 Apr 1;95(7):2434-9.

Related Products

11-4811 Anti-Rat IgG FITC

14-4031 Rat IgG2b K Isotype Control Purified

16-0401 Anti-Mouse CD40 Functional Grade Purified (1C10)

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com