


Anti-Human IL-17A Purified

Catalog Number: 14-7178

Also Known As: Interleukin-17A, IL17A

RUO: For Research Use Only



Immunoprecipitation of IL-17A/F heterodimer with anti-IL-17A clone eBio64CAP17 followed by immunoblot with anti-IL-17F clone H17F10A7. PBMC (1 and 4), in vitro-polarized Th17 (2 and 5), or PMA and Ionomycin-restimulated Th17 (3 and 6) were loaded. Direct lysates (lanes 1-3) are on the left and IP samples (lanes 4-6) are on the right.

Product Information

Contents: Anti-Human IL-17A Purified


REF Catalog Number: 14-7178

Clone: eBio64CAP17


Concentration: 0.5 mg/ml


Host/Isotype: Mouse IgG1, κ

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 Temperature Limitation: Store at 2-8°C.

LOT Batch Code: Refer to Vial

 Use By: Refer to Vial

 Caution, contains Azide

Description

The eBio64CAP17 antibody reacts with human IL-17A; the antibody has been reported to cross react with Rhesus monkey IL-17A, as verified by intracellular staining experiments. The eBio64CAP17 antibody is a neutralizing antibody. Reactivity of the eBio64CAP17 antibody with other IL-17 family members has not been evaluated. Interleukin-17A (IL-17A) is a CD4+ T cell-derived cytokine that promotes inflammatory responses in cell lines and is elevated in rheumatoid arthritis, asthma, multiple sclerosis, psoriasis, and transplant rejection. The cDNA encoding human IL-17A was isolated from a library of CD4+ T cells; the encoded protein exhibits 72 percent amino acid identity with HVS13, an open reading frame from a T lymphotropic Herpesvirus saimiri, and 63 percent with mouse CTLA-8 (cytotoxic T-lymphocyte associated antigen-8). Human IL-17A exists as glycosylated 20-30 kD homodimers. High levels of IL-17A homodimer are produced by activated peripheral blood CD4+ T-cells. IL-17A enhances expression of the intracellular adhesion molecule-1 (ICAM-1) in human fibroblasts. Human IL-17A also stimulates epithelial, endothelial, or fibroblastic cells to secrete IL-6, IL-8, G-CSF, and PGE2. In the presence of human IL-17A, fibroblasts can sustain the proliferation of CD34+ hematopoietic progenitors and induce maturation into neutrophils. Mouse, rat, and human IL-17A can induce IL-6 secretion in mouse stromal cells, indicating that all homologs can recognize the mouse IL-17A receptor.

IL-23-dependent, IL-17A-producing CD4+ T cells (Th-17 cells) have been identified as a unique subset of Th cells that develops along a pathway that is distinct from the Th1- and Th2- cell differentiation pathways. The hallmark effector molecules of Th1 and Th2 cells, e.g., IFN- γ and IL-4, have each been found to negatively regulate the generation of these Th-17 cells.

Additionally, activated human CD4+ T cells have been found to produce the IL-17A/F heterodimer, as well as the corresponding homodimers. In comparing the relative potency of IL-17A, IL-17F, and IL-17A/F, all three were found to induce GRO- α secretion; IL-17A was most potent, followed by IL-17A/F heterodimer, then IL-17F (100fold lower than IL-17A). eBio64CAP17 can be used to detect IL-17 heterodimers by immunoprecipitation followed by immunoblot with H17F10A7 anti-IL17F monoclonal antibody.

The eBio64CAP17 has been shown to react to rhesus and marmoset primates.

Applications Reported

This eBio64CAP17 antibody has been reported for use in intracellular staining followed by flow cytometric analysis, immunoprecipitation, IHC frozen, ELISA, and ELISPOT. (Fluorochrome conjugated eBio64CAP17 is recommended for use in intracellular flow cytometry.)

Applications Tested

The Affinity Purified eBio64CAP17 antibody has been tested as the capture antibody in a sandwich ELISA for measurement of human IL-17A protein levels, in combination with the biotinylated eBio64DEC17 antibody (13-7179) for detection and recombinant human IL-17A (14-8179) as the standard. A suitable range of concentrations of this antibody for ELISA capture is 0.5 – 2.0 $\mu\text{g/ml}$. A standard curve consisting of

doubling dilutions of the recombinant standard over the range of 1000 pg/ml - 8 pg/ml should be included in each ELISA plate.

The Functional Grade Purified eBio64CAP17 antibody has been tested by LAL assay to verify lowest endotoxin levels and has been tested for neutralization of IL-17A bioactivity and for ELISPOT capture.

It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Kap YS, van Meurs M, van Driel N, Koopman G, Melief MJ, Brok HP, Laman JD, 't Hart BA. A monoclonal antibody selection for immunohistochemical examination of lymphoid tissues from non-human primates. *J Histochem Cytochem.* 2009 Dec;57(12):1159-67. (eBio64CAP17, IHC frozen, PubMed)

Related Products

13-7179 Anti-Human IL-17A Biotin (eBio64DEC17)

14-8179 Human IL-17A Recombinant Protein

88-7876 Human IL-17A ELISPOT Ready-SET-Go!®

88-7976 Human IL-17A (Interleukin-17A, IL17A) ELISA Ready-SET-Go! Kit (See replacement item BMS2017)

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com