

# Thermo Scientific Verso 1-Step RT-PCR Kit

## **Description**

Verso<sup>TM</sup> 1-Step RT-PCR Kit supplies in only two vials all the components required to perform a rapid, sensitive and reproducible RT-PCR for the detection and analysis of RNA.

- Verso<sup>TM</sup> Enzyme Mix includes Verso<sup>TM</sup> Reverse Transcriptase, which is active at high temperatures, is highly sensitive and can generate long cDNA strands. This mix also contains RNase inhibitor to protect RNA templates from degradation.
- <u>1-Step PCR Master Mix</u> (2X), a proprietary reaction buffer which has been optimized to allow both reverse transcription and PCR amplification to occur in the same reaction across a wide range of templates.
  - Thermoprime Plus, a DNA Polymerase stable at high temperature.
- <u>RT Enhancer</u> is included to remove contaminating DNA, eliminating the need for DNase I treatment.

## **Kit Contents**

Vial	Pack Size (cap color)	
	A	В
Verso Enzyme Mix	40μl (black)	200µl (black)
1-Step PCR Master Mix (2X)	1ml (red)	5 x 1ml (red)
RT Enhancer	100μl (yellow)	500µl (yellow)

1



## INFORMATION

## Verso<sup>TM</sup> Reverse Transcriptase

Verso<sup>TM</sup> is an RNA-dependent DNA polymerase with a significantly attenuated RNase H activity compared to *Reverse*-iT<sup>TM</sup>. Verso<sup>TM</sup> can synthesize long cDNA strands, up to 11 kb, at a temperature range of 42°C to 57°C. Verso<sup>TM</sup> can reverse transcribe total RNA from 1 pg - 1  $\mu$ g. The recommended amount of total RNA template to use in 1-step kits is between 1 pg - 100 ng.

## Thermoprime Plus DNA Polymerase

Thermoprime Plus has 5' to 3' polymerization and exonuclease activity but lacks 3' to 5' exonuclease activity (proofreading).

#### **RT Enhancer**

RT Enhancer is included to remove contaminating DNA, eliminating the need for DNase I treatment. It degrades double stranded DNA during the transcription of RNA and is inactivated after 2 minutes at 95°C.

#### **Storage Conditions**

Store at -20°C until ready for use. Verso<sup>TM</sup> 1-Step RT-PCR Kit is stable for a minimum of 12 months. Avoid repeated freeze thawing. Shipped on ice within the UK and on dry ice for international and within the US.

#### **Additional Info**

- The use of disposable gloves, RNase and DNase free filter tips and plastics is recommended.
- If DNase I treatment has been performed, RT Enhancer is not required.

## Tips before use

Thaw the reagents on ice. Mix and spin down the solutions before use to recover the maximum amount. Do not vortex the Verso Enzyme Mix or the 1-Step PCR Master Mix.

Briefly centrifuge to avoid bubbles within the wells. Always include a no template control (NTC) and a no enzyme control (NEC).



## PROTOCOL

Example of reaction mix preparation.

The volume of each component is for a 50 µl final reaction.

		Volume	Final Concentration
	Verso Enzyme Mix	1 μl	
	1-Step PCR Master Mix (2X)	25 µl	1X
Reaction	Forward primer (10 µM) <sup>a</sup>	1 μl	200 nM
Mix	Reverse primer (10 μM) <sup>a</sup>	1 μl	200 nM
	RT Enhancer	2.5 µl	
	Water (PCR grade) b	variable	
	Template (RNA) <sup>c</sup>	1 - 5 μl	1 ng
	Total volume	50 µl	

## Example of a **1-Step RT-PCR thermal cycling program**:

	Temp.	Time	Number of cycle
cDNA Synthesis <sup>d</sup>	50°C	15 min	1 cycle
Verso inactivation	95°C	2 min	1 cycle
Denaturation	95°C	20 sec	35 - 45 cycles
Annealing <sup>e</sup>	50-60°C	30 sec	
Extension f	72°C	1 min	
Final extension	72°C	5 min	1 cycle

## **Notes**

- a For optimization, a primer titration should be performed from 50 nM to 500 nM final concentration. Scale up or down the volume and concentration as appropriate.
- b The volume of the total reaction should be completed up to 50  $\mu l$  with water.
- c The amount of total RNA added as a template should be between 1 pg and 100 ng.
- d –Depending on the length of template and degree of secondary structure, the efficiency of the first strand synthesis maybe improved by optimizing temperature and time (42-57°C for 5-30 minutes).
- e Annealing temperature depends on primer sequence.
- f Time of extension depends on the length of the amplicon. If the amplicon exceeds 1 kb amplification time should be adapted (Thermoprime Plus DNA Polymerase extends at approximately 1 kb/min).



## **Quality control**

 $Verso^{TM} \ 1\text{-Step RT-PCR Kit is tested functionally for use in RT-PCR.} \\$ 

## **Ordering Information**

AB-1454/A	Verso™ 1-Step RT-PCR Kit	40 x 50 μl rxns
AB-1454/B	Verso <sup>TM</sup> 1-Step RT-PCR Kit	200 x 50 µl rxns

## **Related Products**

Cat. No.	Description	Quantity
AB-2400	ABgene® SuperPlate™ Semi-Skirted 96-Well PCR Plate	25 plates
AB-2400/W	ABgene® SuperPlate™ Semi-Skirted 96-Well PCR Plate, white	25 plates
AB-2800	ABgene® SuperPlate™ Skirted 96-Well PCR Plate	25 plates
AB-2800/W	ABgene® SuperPlate™ Skirted 96-Well PCR Plate, White	25 plates
AB-0745	Easy Peel (heat seal)	100 sheets
AB-0626	Adhesive PCR Foil	100 sheets
AB-0558	Adhesive PCR Film	100 sheets
AB-0386	Strips of 8 Domed Caps	120 strips
AB-0783	Strips of 8 Flat Caps	120 strips

For technical information or troubleshooting contact Thermo Scientific Genomics Tech Support:

Troubleshooting:	Email	Phone
North America (US, Canada, Central/South America)	Techservice.genomics @thermofisher.com	+1 (800) 235-9880
Europe (EU, Middle East, Africa)	Techservice.emea.genomics @thermofisher.com	(+) 44 1372 840410
Other Countries	www.thermoscientific.com/ dharmacondistributors	

Use of this product is covered by US Patent No. 6,127,155. The purchase of this product includes a limited, non-transferable immunity from suit under the foregoing patent claims for using only this amount of product for the purchaser's own internal research. No right under any other patent claim, no right to perform any patented method and no right to perform commercial services of any kind, including without limitation reporting the results of purchaser's activities for a fee or other commercial consideration, is conveyed expressly, by implication, or by estoppel. This product is for research use only. Diagnostic uses under Roche patents require a separate license from Roche. Further information on purchasing licenses may be obtained by contacting the Director of Licensing, Applied Biosystems, 850 Lincoln Centre Drive, Foster City, California

Revised April, 2011. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.

Literature Code: AB-1454-v8-0411