

Anti-Mouse/Rat CD184 (CXCR4) Purified


Catalog Number: 14-7990

Also Known As: Fusin, CD184

RUO: For Research Use Only

Product Information

Contents: Anti-Mouse/Rat CD184 (CXCR4) Purified

 Catalog Number: 14-7990

Clone: Polyclonal

Host/Isotype: Rabbit IgG


Formulation: Contains 0.09% sodium azide.

The 50 µg size of this antibody is supplied in liquid form. The 200 µg size of this antibody may be supplied in liquid or lyophilized powder forms. To reconstitute the lyophilized antibody, add 200 µL of PBS for a final concentration of 1mg/mL.

 Temperature Limitation: Store at 2-8°C.

 Batch Code: Refer to Vial

 Use By: Refer to Vial

 Caution, contains Azide

Description

The CXCR4 polyclonal antibody reacts with both rat and mouse CXCR4 and was generated using *E. coli*-expressed rat CXCR4 as an immunogen. CXCR4 (Fusin) is a member of the seven-transmembrane domain G-protein-coupled chemokine receptor family. The full-length cDNA was isolated from a human spleen cDNA library. Its ligand is chemokine stroma-derived factor (SDF). CXCR4 has been shown to be the co-receptor for HIV-1 binding to CD4 via the viral envelope glycoprotein gp120. In other cases, CXCR4 can function as the only receptor for the interaction of HIV-2 with CD4 and host cells. A mouse monoclonal antibody, 12G5 to CXCR4 has been shown to inhibit HIV infectivity and HIV-induced syncytium.

Applications Reported

This polyclonal antibody has been reported for use in immunoblotting (WB). Cross-reactivity to CXCR4 in other species has not been determined.

Applications Tested

The tested titer for Western blot is 1:2000. It is recommended that the reagent be carefully titrated for optimal performance in the assay of interest.

References

Ruran Mo, R., et al. 2003. T Cell Chemokine Receptor Expression in Aging. *J. Immunol.* 170: 895-904 [Cites use of the antibody in Western blotting.]

General References:

Federspiel, B. et al. 1993. Molecular cloning of the cDNA and chromosomal localization of the gene for a putative seven-transmembrane segment (7-TMS) receptor isolated from human spleen. *Genomics* 16 (3): 707-12

Nagasawa, T. et al. 1998. A novel CXC chemokine PBSF/SDF-1 and its receptor CXCR4: their functions in development, hematopoiesis and HIV. *Semin Immunol* 10 (3): 179-85

Mcknight, A. et al. 1997. Inhibition of Human Immunodeficiency Virus Fusion by a Monoclonal Antibody to a Coreceptor (CXCR4) Is both Cell Type and Virus Strain Dependent. *J. Virol.*, 71 (2): 1692-1696

Related Products

18-8816 Rabbit TrueBlot®: Anti-Rabbit IgG HRP

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com