



Protease Inhibitor Cocktails

<u>Code</u>	<u>Description</u>	<u>Size</u>
M221-1ml	Protease Inhibitor Cocktail, General Use Supplied as lyophilized powder. <i>Each vial can be reconstituted in 1 ml deionized water to form a 100X solution.</i>	1.0 ml
M222-1ml	Protease Inhibitor Cocktail, General Use with EDTA Supplied as lyophilized powder. <i>Each vial can be reconstituted in 1 ml deionized water to form a 100X solution.</i>	1.0 ml
M250-1ml	Protease Inhibitor Cocktail, Mammalian Supplied as lyophilized powder. <i>Each vial can be reconstituted in 1 ml deionized water to form a 100X solution.</i>	1.0 ml
M306-5ml	Protease Inhibitor Cocktail, Bacterial Supplied as lyophilized powder. <i>Each vial contains sufficient material to provide 5 ml of a 20X solution upon reconstitution.</i>	5.0 ml
M307-1ml	Protease Inhibitor Cocktail, Plant Supplied as lyophilized powder. <i>Each vial can be reconstituted in 1 ml dimethyl sulfoxide to form a 100X solution.</i>	1.0 ml

General Information:

Protease inhibitors are critical reagents for the preservation of protein integrity during purification and analysis procedures. Inhibitor cocktails are extensively used to provide broad spectrum protection during cell lysis and tissue dissolution. These cocktails target a range of different proteases including serine, cysteine, aspartic and metalloproteases as well as aminopeptidases.

AMRESKO offers four protease inhibitor cocktails designed to preserve the protein composition of samples in a broad range of organisms. A general use cocktail is available with or without EDTA, while specific formulations are available for mammalian, bacterial or plant lysates. All formulations are prepared from ultrapure reagents to ensure maximum inhibition of the specified proteases. Cocktails are provided as convenient, easy-to-use powders that can be reconstituted in the supplied vials.

Check the current AMRESKO catalog for a comprehensive listing of all AMRESKO protease inhibitors.

Storage/Stability:

The lyophilized powder should be stored at -20°C where it is stable for up to 1 year. The reconstituted solution should be stored at -20°C where it is stable for up to 2 weeks. Repeated freezing and thawing cycles of the reconstituted solution is not recommended.

Application Disclaimer

*For Research Use Only.
Not for Therapeutic or Diagnostic Use.*

Protocols and Formulations

➔ **Note:** Reconstituted cocktails should be aliquoted into multiple tubes and stored at -20°C. Multiple freezing and thawing cycles are not recommended.

M221-1ML, General Use Protease Inhibitor Cocktail
M222-1ML, General Use Protease Inhibitor Cocktail with EDTA

Preparation of 100X Solution:

1. Add 1 ml deionized, distilled water and mix gently until powder is completely resuspended.
2. Aliquot into multiple tubes and store at -20°C.

<u>Name</u>	<u>MW</u>	<u>100X Concentration</u>	<u>Recommended Working Concentration</u>
AEBSF	239.5	50.00 mM	0.50 mM
Aprotinin	6512.0	30.00 uM	0.30 uM
Bestatin	308.4	1.00 mM	10.00 uM
E-64	357.4	1.00 mM	10.00 uM
Leupeptin	493.6	1.00 mM	10.00 uM
EDTA*	372.2	5.00 mM	50.00 uM

* Present only in M222- 1ML

M250-1ML, Mammalian Protease Inhibitor Cocktail:

Preparation of 100X Solution:

1. Add 1 ml deionized, distilled water and mix gently until powder is completely resuspended.
2. Aliquot into multiple tubes and store at -20°C.

<u>Name</u>	<u>MW</u>	<u>100X Concentration</u>	<u>Recommended Working Concentration</u>
AEBSF	239.5	120.00 mM	1.20 mM
Aprotinin	6512.0	46.00 uM	0.46 uM
Bestatin	308.4	1.36 mM	14.00 uM
E-64	357.4	1.23 mM	12.30 uM
Leupeptin	493.6	11.20 mM	112.00 uM
Pepstatin	685.9	116.00 uM	1.16 uM

M306-5ML Bacterial Protease Inhibitor Cocktail

Preparation of 20X Solution:

1. Add 1 ml DMSO to vial and vortex 1 minute.
2. Add 4 ml deionized, distilled water and mix.
3. Aliquot into multiple tubes and store at -20°C. (solution may be cloudy and precipitate may be present)

<u>Name</u>	<u>MW</u>	<u>20X Concentration</u>	<u>Recommended Working Concentration*</u>
AEBSF	239.5	22.00 mM	1.10 mM
Bestatin	308.4	1.87 mM	9.35 uM
E-64	357.4	220.00 uM	11.00 uM
Pepstatin	685.9	2.20 mM	110.00 uM
EDTA	372.2	93.00 mM	4.65 mM

*1 ml of solution is sufficient to inhibit 20 ml of lysate from 4 g (wet weight) of *E. coli* cells.

M307-1ML, Plant Protease Inhibitor Cocktail:

Preparation of 100X Solution:

1. Add 1 ml dimethyl sulfoxide and mix gently until powder is completely resuspended.
2. Aliquot into multiple tubes and store at -20°C.

<u>Name</u>	<u>MW</u>	<u>100X Concentration</u>	<u>Recommended Working Concentration</u>
AEBSF	239.5	220.00 mM	2.20 mM
Bestatin	308.4	10.70 mM	107.00 uM
E-64	357.4	3.00 mM	30.00 uM
Leupeptin	493.6	2.00 mM	20.00 uM
Pepstatin	685.9	2.20 mM	22.00 uM
1,10-Phenanthroline	198.2	550.00 mM	5.50 mM

Inhibitor Specificity

Name	Inhibition Specificity
AEBSF	<ul style="list-style-type: none"> Irreversible serine protease inhibitor <ul style="list-style-type: none"> Chymotrypsin Trypsin Kallikrein Plasmin Thrombin. Non-toxic alternative for PMSF
Aprotinin	<ul style="list-style-type: none"> Competitive, reversible serine protease inhibitor <ul style="list-style-type: none"> Chymotrypsin Trypsin Kallikrein Plasmin Does not inhibit Factor Xa or thrombin
Bestatin	<ul style="list-style-type: none"> Competitive aminopeptidases inhibitor <ul style="list-style-type: none"> Aminopeptidase B Leucine aminopeptidase Tripeptide aminopeptidase Does not inhibit carboxypeptidases
E-64	<ul style="list-style-type: none"> Irreversible cysteine protease inhibitor <ul style="list-style-type: none"> Papain Calpain Cathepsin B, H, L and S Effective inhibitor of collagenase
EDTA	<ul style="list-style-type: none"> Reversible metalloprotease inhibitor – Chelates metal ions
Leupeptin	<ul style="list-style-type: none"> Reversible cysteine and serine protease inhibitor <ul style="list-style-type: none"> Trypsin Plasmin Papain Kallikrein Thrombin Cathepsin A and B Effective inhibitor of collagenase
Pepstatin	<ul style="list-style-type: none"> Aspartic acid protease inhibitor <ul style="list-style-type: none"> Pepsin Renin Cathepsin D Insoluble in water
1,10-Phenanthroline	<ul style="list-style-type: none"> Metalloprotease inhibitor – Chelates iron and other divalent cations

Related Products
Code
Peroxide Free Detergents

M228-10ML-5PK

M236-10ML-5PK

Detergents

0777

0694

E109

0479

Buffers

0826

0780

Reagents for Protein Purification

E177

0105-500G

0281-5G

Product

Tween® 20, 10% Solution

Triton®X-100, 10% Solution

Tween® 20

Triton® X-100

Nonidet® P-40 Substitute

n-Octyl-βD-glucopyranoside

Tris

Phosphate Buffered Saline

EDTA Disodium Salt Dihydrate

DTT (DL-Dithiothreitol)

ZY0276 Rev. 7/9/12 ©Copyright 2006, 2007 by AMRESCO Inc. All Rights Reserved. AMRESCO® is a registered trademark of AMRESCO Inc.



Corporate Headquarters

AMRESCO LLC.
6681 Cochran Road
Solon, OH 44139 USA
Tel: (440) 349-1199
Toll free: 1-800-448-4442 (US & Canada)
Fax: (440) 349-2441
e-mail: info@amresco-inc.com

To Order (USA & Canada)

www.amresco-inc.com
Toll free: 1-800-448-4442
Toll free fax: 1-800-326-3733
Fax: (440) 349-3255
e-mail: dom-sales@amresco-inc.com

International Orders:

Tel: (440) 349-1199
Fax: (440) 349-2441
e-mail: int-sales@amresco-inc.com

Technical Support

Toll free: 1-800-610-2789 (USA & Canada)
Fax: (440) 349-0235
e-mail: techinquiry@amresco-inc.com

www.amresco-inc.com