


Anti-Human CD284 (TLR4) Purified

Catalog Number: 14-9917

Also Known As: TLR-4, toll-like receptor 4

RUO: For Research Use Only



Surface staining of normal human peripheral blood cells with Anti-Human CD284 (TLR4) PE (left), and Anti-Human CD284 (TLR4) APC (right). Appropriate isotype controls were used (open histogram). Cells in the monocyte population were used for analysis.

Product Information

Contents: Anti-Human CD284 (TLR4) Purified


REF Catalog Number: 14-9917

Clone: HTA125


Concentration: 0.5 mg/ml


Host/Isotype: Mouse IgG2a, κ

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 Temperature Limitation: Store at 2-8°C.

LOT Batch Code: Refer to Vial

 Use By: Refer to Vial

 Caution, contains Azide

Description

The HTA125 monoclonal antibody reacts with human Toll-like receptor 4 (TLR4). So far, at least ten members of the Toll family have been identified in humans. This family of type I transmembrane proteins is characterized by an extracellular domain with leucine-rich repeats and a cytoplasmic domain with homology to the type I IL-1 receptor. Two of these receptors, TLR2 and TLR4, are pattern recognition receptors and signaling molecules in response to bacterial lipoproteins and have been implicated in innate immunity and inflammation. TLR4 physically associates with another molecule called MD-2, and together with CD14, this complex is responsible for LPS recognition and signaling. TLR4 is expressed by peripheral blood monocytes. HTA125 has been reported to immunoprecipitate human TLR4 (~100 kDa) from transfected cells. Most TLR cell surface expression, especially TLR1 and TLR4, occurs at low levels on monocytes and at even lower levels on other cell types including granulocytes and immature dendritic cells (iDC). Furthermore, a relatively high degree of variability in TLR surface expression has been reported among normal donors.

Applications Reported

The HTA125 antibody has been reported for use in flow cytometric analysis, and immunoprecipitation. It has also been reported in blocking of LPS-induced cytokine production. For detection of peripheral monocytes, a three step staining protocol is recommended using purified anti-human TLR4 followed by biotin anti-mouse IgG and streptavidin-PE. (Please use Functional Grade purified HTA125, cat. 16-9917, in functional assays.)

Applications Tested

The HTA125 antibody has been tested by flow cytometric analysis of human peripheral blood leukocytes. This can be used at less than or equal to 1 μ g per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

It is highly recommended that for optimal staining of TLR4, whole blood be stained using the LWB protocol rather than Ficoll-gradient prepared PBMCs. The use of a density gradient appears to reduce the staining intensity significantly.

References

Tabeta, K., K. Yamazaki, et al. 2000. Toll-like receptors confer responsiveness to lipopolysaccharide from *Porphyromonas gingivalis* in human gingival fibroblasts. *Infect Immun.* 68(6): 3731-5.

Akashi, S., H. Ogata, et al. 2000. Regulatory roles for CD14 and phosphatidylinositol in the signaling via toll-like receptor 4-MD-2. *Biochem Biophys Res Commun.* 268(1): 172-7.

Shimazu, R., S. Akashi, et al. 1999. MD-2, a molecule that confers lipopolysaccharide responsiveness on Toll-like receptor 4. *J Exp Med.* 189(11):

1777-82.

de Graaf R, Kloppenburg G, et al. 2006. Human heat shock protein 60 stimulates vascular smooth muscle cell proliferation through Toll-like receptors 2 and 4. *Microbes Infect.* 8(7):1859-65. (HTA125, IHC paraffin, PubMed)

Mirlashari MR and Lyberg T. 2003. Expression and involvement of Toll-like receptors (TLR)2, TLR4, and CD14 in monocyte TNF-alpha production induced by lipopolysaccharides from *Neisseria meningitidis*. *Med Sci Monit.* 9(8):BR316-24. (HTA125, FA, PubMed)

Related Products

12-4317 Streptavidin PE

13-4013 Anti-Mouse IgG Biotin (Polyclonal)

14-4724 Mouse IgG2a K Isotype Control Purified

14-8185 B18R Recombinant Protein

14-9920 Anti-Human CD284 (TLR4) Purified (Polyclonal)

34-8185 B18R Recombinant Protein Carrier-Free

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com