

Anti-Mouse CD273 (B7-DC) Purified

Catalog Number: 14-9972

Also Known As: B7DC, PD-L2, PDL-2, PDL2

RUO: For Research Use Only

Product Information

Contents: Anti-Mouse CD273 (B7-DC) Purified


 Catalog Number: 14-9972

Clone: 122


Concentration: 0.5 mg/ml


Host/Isotype: Rat IgG2a, κ

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 Temperature Limitation: Store at 2-8°C.

 Batch Code: Refer to Vial

 Use By: Refer to Vial

 Caution, contains Azide

Description

The 122 monoclonal antibody reacts with mouse B7-DC, also known as PD-L2. B7-DC, a recently identified member of the B7 family, has a predicted molecular weight of approximately 25 kDa and belongs to the Ig superfamily. The mouse B7-DC has a short cytoplasmic tail (4aa). B7-DC is primarily expressed by subpopulations of dendritic cells and monocytes/macrophages in the mouse. Although B7-DC has structural and sequence similarities to the B7 family, it does not bind CD28/CTLA-4, rather it is a ligand for PD-1. The interactions between PD-1 and B7-DC/PD-L2 have been reported to be involved in costimulation or suppression of T cell proliferation depending on state of cellular activation. 122 has been demonstrated to block binding of TY25 (Cat. No. 14-5986), another mAb specific for mouse B7-DC.

Applications Reported

122 has been reported for use in flow cytometric analysis.

Applications Tested

The 122 antibody has been tested by flow cytometric analysis of mouse dendritic cell suspensions and mouse B7-DC transfected cells. This can be used at less than or equal to 0.5 μ g per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Ishida M, Iwai Y, Tanaka Y, Okazaki T, Freeman GJ, Minato N, Honjo T. 2002. Differential expression of PD-L1 and PD-L2, ligands for an inhibitory receptor PD-1, in the cells of lymphohematopoietic tissues. *Immunol Lett.* 84(1):57-62. Carter L, Fouser LA, Jussif J, Fitz L, Deng B, Wood CR, Collins M, Honjo T, Freeman GJ, Carreno BM. 2002. PD-1:PD-L inhibitory pathway affects both CD4(+) and CD8(+) T cells and is overcome by IL-2. *Eur J Immunol.* 32(3):634-43.

Related Products

11-4317 Streptavidin FITC

11-4811 Anti-Rat IgG FITC

12-4317 Streptavidin PE

12-5986 Anti-Mouse CD273 (B7-DC) PE (TY25)

13-4813 Anti-Rat IgG Biotin (Polyclonal)

13-5986 Anti-Mouse CD273 (B7-DC) Biotin (TY25)

14-4321 Rat IgG2a K Isotype Control Purified

14-5986 Anti-Mouse CD273 (B7-DC) Purified (TY25)

16-5986 Anti-Mouse CD273 (B7-DC) Functional Grade Purified (TY25)

17-4317 Streptavidin APC

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com