

Product Data Sheet

PE anti-mouse CD4

Catalog # / Size: 130310 / 200 µg

Clone: H129.19

Isotype: Rat IgG2a, κ

Immunogen: Mouse lymphocytes


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



C57BL/6 splenocytes stained with H129.19 PE

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: blocking of CD4⁺ T cell activation¹⁻³, depletion. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays.

Application References:

1. Naquet P. 1985. *Immunogenetics*. 22:247-256.
2. Pierres A. 1984. *J. Immunol.* 132:2775-2782.
3. Guo MW. 1995. *Zygote*. 3:65-73.
4. Ghobrial RR. 1989. *Clin Immunol Immunopathol*. 52:486-506.

Description: CD4 is a 55 kD protein, also known as L3T4 or T4. It is a member of the Ig superfamily, primarily expressed on most thymocytes, a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a coreceptor with the TCR during T cell activation and thymic differentiation by binding MHC class II and associating with the protein tyrosin kinase, lck. The H129.19 antibody is able to block CD4 mediated cell adhesion and T cell activation. It is also useful for in vivo and in vitro depletion of CD4⁺ T cells. H129.19 antibody blocks binding of the anti-mouse CD4 mAbs GK1.5 and RM4-5, but not RM4-4 antibody. H129.19 antibody inhibits various responses of T helper cells to antigenic or mitogenic stimuli.

Antigen References:

1. Barclay A, et al. 1997. *The Leukocyte Antigen FactsBook Academic Press*.
2. Bierer BE, et al. 1989. *Annu. Rev. Immunol*. 7:579.
3. Janeway CA. 1992. *Annu. Rev. Immunol*. 10:645.
4. Naquet P. 1985. *Immunogenetics*. 22:247-256.
5. Pierres A. 1984. *J. Immunol.* 132:2775-2782.
6. Guo MW. 1995. *Zygote*. 3:65-73.
7. Ghobrial RR. 1989. *Clin Immunol Immunopathol*. 52:486-506.

Related Products:	Product	Clone	Application
	PE Rat IgG2a, κ Isotype Ctrl	RTK2758	FC, ICFC
	Cell Staining Buffer		FC, ICC, ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	TruStain fcX™ (anti-mouse CD16/32)	93	FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.