

Product Data Sheet

PE anti-mouse CD9

Catalog # / Size: 124805 / 50 μg

124806 / 200 µg

Clone: MZ3

Isotype: Rat IgG2a, κ **Reactivity:** Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with

PE under optimal conditions. The solution is free of unconjugated PE and


unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



C57BL/6 bone marrow cells stained with RB6-8C5 (Gr-1) PerCP/Cy5.5 and MZ3 PE

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For

immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is

recommended that the reagent be titrated for optimal performance for each application.

Application References: 1. Won WJ, et al. 2002. J. Immunol. 168:5605. (FC, WB)

Description: CD9 is a surface glycoprotein of the tetraspannin family. It is expressed on a variety of cells, including nerve, muscle

cells and many cells of hematopoietic origin. CD9 is found to participate in forming a large molecular cell complex with other member proteins, such as MHC class II, CD19, CD5 and other TM4SF molecules. It is reported that CD9 is a marker of marginal zone B cells, B1 cells and plasma cells. The diverse functions of CD9 may largely depend upon its

associated molecules on different cells.

Antigen References: 1. Boucheix C, et al. 1991. J. Biol. Chem. 266:117

2. Lanza F, et al. 1991. J. Biol. Chem. 266:10638

RBC Lysis Buffer (10X)

TruStain fcX™ (anti-mouse CD16/32)

FC, ICFC

FC

FC

FC



