

Product Data Sheet

FITC anti-mouse CD9

Catalog # / Size: 124807 / 25 µg

124808 / 100 µg

Clone: MZ3

Isotype: Rat IgG2a, κ

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with

FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent

staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is $\le 1.0 \ \mu g$ per 10^6 cells in 100 μl volume. It is recommended that the reagent be titrated for optimal performance for each

application.

Application References: 1. Won WJ, et al. 2002. J. Immunol. 168:5605. (FC, WB)

Description: CD9 is a surface glycoprotein of the tetraspannin family. It is expressed on a variety of cells, including nerve, muscle cells and many cells of hematopoietic origin. CD9 is found to participate in forming a large molecular cell complex with

other member proteins, such as MHC class II, CD19, CD5 and other TM4SF molecules. It is reported that CD9 is a marker of marginal zone B cells, B1 cells and plasma cells. The diverse functions of CD9 may largely depend upon its

associated molecules on different cells.


Antigen References: 1. Boucheix C, et al. 1991. J. Biol. Chem. 266:117

2. Lanza F, et al. 1991. J. Biol. Chem. 266:10638

Application Related Products: Product Clone RTK2758 FITC Rat IgG2a, κ Isotype Ctrl

FC, ICFC FC, ICFC Cell Staining Buffer RBC Lysis Buffer (10X) 93

TruStain fcX™ (anti-mouse CD16/32)



C57BL/6 bone marrow cells stained with RB6-8C5 (Gr-1) PerCP/Cy5.5 and MZ3 FITC



