

Product Data Sheet

Purified anti-mouse CD8b.2

Catalog # / Size: 140401 / 50 µg
 140402 / 500 µg

Clone: 53-5.8

Isotype: Rat IgG1, κ

Immunogen: Mouse thymus or spleen


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



C57BL/6 mouse splenocytes stained with purified CD8b.2 (53-5.8) conjugated with FITC and CD8a (53-6.7) APC

Applications:

Applications: FC - *Quality tested*
 IF, IHC, IP - *Reported in the literature*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunofluorescence³, immunohistochemical staining⁴ of frozen tissue section using dry ice-isopentane method and immunoprecipitation⁵.

Application References:

1. Ledbetter J, et al. 1979. *Immunol. Rev.* 47:63.
2. Ledbetter J, et al. 1980. *J. Exp. Med.* 152:280.
3. Vremec D, et al. 2000. *J. Immunol.* 164:2978. (IF)
4. Lawrence D, et al. 1999. *J. Virol.* 73:1795. (IHC)
5. Bosselut R, et al. 1999. *J. Exp. Med.* 190:1517. (IP)

Description: CD8b is the 32 kD β chain of CD8, also known as Lyt-3.2 or Ly-3.2. It is a member of the Ig superfamily expressed as a heterodimer with the CD8α chain on a subset of MHC class I-restricted T cells and most thymocytes. CD8 is a co-receptor for the TCR complex involved in T cell activation. The antibody 53-5.8 is specific for Ly-3.2 and has low reactivity with Ly-3.1.

Antigen References:

1. Ledbetter J, et al. 1981. *J. Exp. Med.* 153:1503.
2. Renard V, et al. 1996. *J. Exp. Med.* 184:2439.

Related Products:

Product	Clone	Application
Purified Rat IgG1, κ Isotype Ctrl	RTK2071	FC, ICFC, ICC, IF, IHC, IP, WB
Cell Staining Buffer		FC, ICC, ICFC
RBC Lysis Buffer (10X)		FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.