


Anti-Human CD16 PE

Catalog Number: 12-0167

Also Known As: Low Affinity IgG Receptor 3, FCGR3a, FCGR3, IGFR3

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of normal human peripheral blood cells with Anti-Human CD56 (NCAM) APC and Mouse IgG1 kappa Isotype Control PE (cat. 12-4714) (left) or Anti-Human CD16 PE (right). Cells in the granulocyte gate were used for analysis.

Product Information

Contents: Anti-Human CD16 PE


REF **Catalog Number:** 12-0167

Clone: B73.1


Concentration: 5 µL (0.5 µg)/test

Host/Isotype: Mouse IgG1, kappa

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 **Temperature Limitation:** Store at 2-8°C. Do not freeze. Light sensitive material.

LOT **Batch Code:** Refer to Vial

 **Use By:** Refer to Vial

 **Caution, contains Azide**

Description

This B73.1 monoclonal antibody reacts with human CD16, a member of the Fc receptor family also known as Fc gamma RIIIa and Fc gamma RIIb. This 50-65 kDa receptor is expressed on natural killer cells, neutrophils, monocytes, and macrophages. Crosslinking of CD16 in natural killer cells and monocytes results in activation of the MAPK and PI3K signaling pathways, thus regulating the production of TNF alpha, IL-6, and IL-1 beta.

This monoclonal antibody has been shown to block Fc receptor functions.

Applications Reported

This B73.1 (B73.1.1) antibody has been reported for use in flow cytometric analysis.

Applications Tested

This B73.1 (B73.1.1) antibody has been pre-titrated and tested by flow cytometric analysis of normal peripheral blood cells. This can be used at 5 µl (0.5 µg)/per test. A test is defined as the amount (µg)/test of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test.

References

Kramer PR, Winger V, Reuben J. PI3K limits TNF-alpha production in CD16-activated monocytes. Eur J Immunol. 2009 Feb;39(2):561-70.

Trotta R, Kanakaraj P, Perussia B. Fc gamma R-dependent mitogen-activated protein kinase activation in leukocytes: a common signal transduction event necessary for expression of TNF-alpha and early activation genes. J Exp Med. 1996 Sep 1;184(3):1027-35.

Perussia B, Starr S, Abraham S, Fanning V, Trinchieri G. Human natural killer cells analyzed by B73.1, a monoclonal antibody blocking Fc receptor functions. I. Characterization of the lymphocyte subset reactive with B73.1. J Immunol. 1983 May;130(5):2133-41. (**B73.1**, FC, FA)

Perussia B, Acuto O, Terhorst C, Faust J, Lazarus R, Fanning V, Trinchieri G. Human natural killer cells analyzed by B73.1, a monoclonal antibody blocking Fc receptor functions. II. Studies of B73.1 antibody-antigen interaction on the lymphocyte membrane. J Immunol. 1983 May;130(5):2142-8. (**B73.1**, FC)

Related Products

12-0168 Anti-Human CD16 PE (eBioCB16 (CB16))
12-4714 Mouse IgG1 K Isotype Control PE (P3.6.2.1)
17-0569 Anti-Human CD56 (NCAM) APC (MEM188 (MEM-188))

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com