
Anti-Mouse CD30 PE


Catalog Number: 12-0301

Also Known As: TNFRSF8, Ki-1

RUO: For Research Use Only

Product Information

Contents: Anti-Mouse CD30 PE

 Catalog Number: 12-0301

Clone: mCD30.1

Concentration: 0.2 mg/ml

Host/Isotype: Armenian Hamster IgG

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer



Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.



Batch Code: Refer to Vial



Use By: Refer to Vial



Caution, contains Azide

Description

The mCD30.1 monoclonal antibody reacts with mouse CD30, a 105-120 kDa member of the TNFR superfamily. CD30 is not detectable on splenocytes, but can be upregulated on T cells stimulated *in vitro* through TCR with a peak expression around day 4-5 post-activation. CD30 is expressed by both CD8⁺ and CD4⁺ activated cells. It is suggested that CD30 regulates cytokine secretion by T cells and has a role in T cell development.

Applications Reported

The mCD30.1 antibody has been reported for use in flow cytometric analysis.

Applications Tested

The mCD30.1 antibody has been tested by flow cytometric analysis of resting and activated mouse splenocyte suspensions. This can be used at less than or equal to 1 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Bowen, M. A., R. K. Lee, et al. (1996). "Structure and expression of murine CD30 and its role in cytokine production." *J Immunol* 156(2): 442-9.

Related Products

12-4888 Armenian Hamster IgG Isotype Control PE (eBio299Arm)

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com