

Product Data Sheet

PE anti-mouse/rat/human CD27

Catalog # / Size: 124209 / 50 µg
124210 / 200 µg

Clone: LG.3A10

Isotype: Armenian Hamster IgG

Immunogen: Armenian hamster fibroblast line ARHO12 transfected with mouse CD27 cDNA


Reactivity: Mouse, Rat, Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



C57BL/6 mouse splenocytes stained with LG.3A10 PE

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application References:

1. Gravestien LA, *et al.* 1995. *Int. Immunol.* 7:551
2. Gravestien LA, *et al.* 1996. *J. Exp. Med.* 184:675
3. Takeda K, *et al.* 2000. *J. Immunol.* 164:1741
4. Welner RS, *et al.* 2009. *J. Immunol.* 183:7768. PubMed

Description: CD27 is also known as S152 and T14. A member of the tumor necrosis factor receptor (TNFR) superfamily, it is a 45 kD protein expressed on peripheral T cells, memory B cells, NK cells, and thymocyte subset. Through its ligand, CD70, CD27 plays a key role in T cell and B cell interactions. Additionally, ligation of CD27 on naïve T cells may be important in their maturation to effector cells.

Antigen References:

1. Hintzen RQ, *et al.* 1994. *Immunol. Today* 15:307.
2. Akiba H, *et al.* 1998. *J. Biol. Chem.* 273:13353.
3. Kobata T, *et al.* 1995. *P. Natl. Acad. Sci. USA* 92:11249.
4. Hendriks J, *et al.* 2000. *Nat. Immunol.* 1:433
5. Hintzen RQ, *et al.* 1994. *Int. Immunol.* 6:477
6. Camerini D, *et al.* 1991. *J. Immunol.* 147:3165

Related Products:

Product
 Cell Staining Buffer
 RBC Lysis Buffer (10X)
 PE Armenian Hamster IgG Isotype Ctrl
 TruStain fcX™ (anti-mouse CD16/32)

Clone

HTK888
93

Application

FC, ICC, ICFC
 FC, ICFC
 FC, ICFC
 FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.