

## PE anti-mouse CD41

**Catalog # / Size:** 133905 / 25 µg  
133906 / 100 µg

**Clone:** MWRReg30

**Isotype:** Rat IgG1, κ

**Immunogen:** Mouse platelets


**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.2 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



C57 BL/6 platelets were stained with CD41 (clone: MWRReg30) PE (filled histogram) or rat IgG1, κ PE isotype control (open histogram).

## Applications:

**Applications:** FC - Quality tested

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** Additional reported applications (for the relevant formats) include: depletion of platelets *in vivo*.

**Application References:**

1. Nieswandt B, *et al.* 1999. *Blood* 94:684.
2. Teeling JL, *et al.* 2001. *Blood* 98:1095.
3. Bertrand JY, *et al.* 2005. *P. Natl. Acad. Sci. USA* 102:134.
4. Nocito A, *et al.* 2007. *Hepatology* 45:369. (Deplete)
6. Sullivan BP, *et al.* 2010. *Toxicol. Sci.* 115:286. (Deplete) PubMed

**Description:** CD41, also known as integrin α2b and GPIIb, is a transmembrane glycoprotein that is expressed by platelets and megakaryocytes. It was reported that CD41 is also expressed on hematopoietic progenitors. CD41 associates with CD61 (integrin β3) to form complexes that interact with fibrinogen, fibronectin, von Willebrand factor, and thrombin. CD41 is required for platelet adhesion and aggregation. Defect of CD41 leads to disorders of coagulation.

**Antigen References:**

1. Bakewell SJ, *et al.* 2003. *P. Natl. Acad. Sci. USA* 100:14205.
2. Phillips DR, *et al.* 1991. *Cell.* 65:359.

### Related Products:

Product	Clone	Application
PE Rat IgG1, κ Isotype Ctrl	RTK2071	FC, ICFC
Cell Staining Buffer		FC, ICC, ICFC
RBC Lysis Buffer (10X)		FC, ICFC
TruStain fcX™ (anti-mouse CD16/32)	93	FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.