


Anti-Human CD86 (B7-2) PE

Catalog Number: 12-0869

Also known as: B72, B70

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of normal human peripheral blood cells with Mouse IgG2b kappa Isotype Control PE (cat. 12-4732) (blue histogram) or Anti-Human CD86 (B7-2) PE (purple histogram). Cells in the monocyte gate were used for analysis.

Product Information



Contents: Anti-Human CD86 (B7-2) PE

Catalog Number: 12-0869

Clone: IT2.2

Concentration: 5 μ L (0.5 μ g)/test

Host/Isotype: Mouse IgG2b, kappa

HLDA Workshop: VI B-CD86.8



Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.



Batch Code: Refer to vial



Use By: Refer to vial



Caution, contains Azide

Description

The IT2.2 monoclonal antibody reacts with human CD86, an ~80 kDa surface receptor also known as B7-2. CD86 and CD80 are members of the B7 family of costimulatory molecules. CD86 is expressed at low levels on B cells, macrophages, and dendritic cells and is upregulated on B cells through a variety of surface stimuli including the BCR complex, CD40 and some cytokine receptors. In addition to CD80 (B7-1), CD86 is a counter-receptor for the T cell surface molecules CD28 and CD152 (CTLA-4). The interaction of CD86 with its ligands plays a critical role in T-B crosstalk, T cell costimulation, autoantibody production and Th2-mediated Ig production. The kinetics of upregulation of CD86 upon stimulation supports its major contribution during the primary phase of an immune response.

Applications Reported

The IT2.2 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This IT2.2 antibody has been pre-titrated and tested by flow cytometric analysis of human peripheral blood leukocytes. This can be used at 5 μ L (0.5 μ g) per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test.

References

Kapsogeorgou EK, Moutsopoulos HM, Manoussakis MN. Functional expression of a costimulatory B7.2 (CD86) protein on human salivary gland epithelial cells that interacts with the CD28 receptor, but has reduced binding to CTLA4. *J Immunol.* 2001 Mar 1;166(5):3107-13. (IT2.2, FA, PubMed)

Kishimoto T, von dem Borne AEG, et al. eds. *Leucocyte Typing VI: White Cell Differentiation Antigens.* 1998. Garland Publishing Inc. London.

Jeannin P, Delneste Y, Lecoanet-Henchoz S, Gauchat JF, Ellis J, Bonnefoy JY. CD86 (B7-2) on human B cells. A functional role in proliferation and selective differentiation into IgE- and IgG4-producing cells. *J Biol Chem.* 1997 Jun

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com

Anti-Human CD86 (B7-2) PE

Catalog Number: 12-0869

Also known as: B72, B70

RUO: For Research Use Only. Not for use in diagnostic procedures.

20;272(25):15613-9. (IT2.2, FA, PubMed)

Azuma M, Ito D, Yagita H, Okumura K, Phillips JH, Lanier LL, Somoza C. B70 antigen is a second ligand for CTLA-4 and CD28. Nature. 1993 Nov 4;366(6450):76-9. (IT2.2, FC, IP, PubMed)

Related Products

12-4732 Mouse IgG2b K Isotype Control PE

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com