


An Affymetrix Company

# Anti-Mouse CD90.2 (Thy-1.2) PE

Catalog Number: 12-0903

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of C57Bl/6 splenocytes with Anti-Mouse CD3e APC (cat. 17-0031) and 0.06 ug of Rat IgG2b kappa Isotype Control PE (cat. 12-4031) (left) or 10.06 ug of Anti-Mouse CD90.2 (Thy-1.2) PE (right). Total viable cells were used for analysis.

### **Product Information**

Contents: Anti-Mouse CD90.2 (Thy-1.2) PE


REF Catalog Number: 12-0903

Clone: 30-H12

Concentration: 0.2 mg/mL Host/Isotype: Rat IgG2b, kappa



**Formulation:** aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer **Temperature Limitation:** Store at 2-8°C. Do not





Use By: Refer to vial



The 30-H12 monoclonal antibody reacts with mouse CD90.2, also known as Thy-1.2, a GPI-linked membrane molecule. CD90.2 is expressed by mouse thymocytes and mature T cells as well as neurons in CD90.2-expressing mouse strains. These strains include BALB/c, CBA, C3H, C57BL/6, C58/, SJL and others. Cells from CD90.1-expressing strains including PL and AKR do not stain with 30-H12. CD90 is involved in regulation of adhesion and signal transduction by T cells.

#### **Applications Reported**

The 30-H12 antibody has been reported for use in flow cytometric analysis.

#### **Applications Tested**

The 30-H12 antibody has been tested by flow cytometric analysis of mouse splenocyte suspensions. This can be used at less than or equal to 0.125  $\mu$ g per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

#### References

Sugai M, Kondo S, Shimizu A, Honjo T. Isolation of differentially expressed genes upon immunoglobulin class switching by a subtractive hybridization method using uracil DNA glycosylase. Nucleic Acids Res. 1998 Feb 5;26(4):911-8. (In vitro depletion)

Ledbetter, J.A. and L.A. Herzenberg (1979). Xenogenic monoclonal antibodies to mouse lymphoid differentiation antigens. Immunol Rev 47: 63-90.

## **Related Products**

12-4031 Rat IgG2b K Isotype Control PE 17-0031 Anti-Mouse CD3e APC (145-2C11)

info@ebioscience.com