


Anti-Mouse CD137 (4-1BB) PE

Catalog Number: 12-1371

Also Known As: Ly-63, TNFRSF9, ILA

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of 3-day unstimulated (left) and 3-day ConA-stimulated (right) BALB/c splenocytes with 0.125 ug of Golden Syrian Hamster IgG Isotype Control PE (cat. 12-4914) (open histogram) or 0.125 ug of Anti-Mouse CD137 (4-1BB) PE (filled histogram). Total viable cells were used for analysis.

Product Information

Contents: Anti-Mouse CD137 (4-1BB) PE


REF Catalog Number: 12-1371

Clone: 17B5

Concentration: 0.2 mg/mL


Host/Isotype: Golden Syrian Hamster IgG

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 **Temperature Limitation:** Store at 2-8°C. Do not freeze. Light sensitive material.

LOT **Batch Code:** Refer to Vial

 **Use By:** Refer to Vial

 **Caution, contains Azide**

Description

The 17B5 monoclonal antibody reacts with mouse CD137, also known as 4-1BB, a member of the TNFR superfamily. CD137 is expressed as monomers, dimers, and tetramers on the surface of activated T cells around day 4-6 after activation through the TCR complex. Interaction of CD137 with its ligand, 4-1BBL, is reported to deliver a costimulatory signal important in the T-APC interaction. Recent data suggests CD137 is expressed on Tregs and may play a role in their proliferation. The monoclonal antibody 17B5 has strong antagonistic effect in *in vitro* functional assays.

Applications Reported

The 17B5 antibody has been reported for use in flow cytometric analysis.

Applications Tested

The 17B5 antibody has been tested by flow cytometric analysis of activated mouse splenocyte cultures. This can be used at less than or equal to 0.25 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Zheng G, Wang B, Chen A. The 4-1BB costimulation augments the proliferation of CD4+CD25+ regulatory T cells. *J Immunol.* 2004 Aug 15;173(4):2428-34. (**17B5**, FA, PubMed)

Related Products

12-4914 Golden Syrian Hamster IgG Isotype Control PE (n/a)

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com